

METRIC FIXED POINT THEORY AND PARTIAL IMPREDICATIVITY

DAVID FERNÁNDEZ-DUQUE, PAUL SHAFER, HENRY TOWNSNER, AND KEITA YOKOYAMA

ABSTRACT. We show that the Priess-Crampe & Ribenboim fixed point theorem is provable in RCA_0 . Furthermore, we show that Caristi's fixed point theorem for both Baire and Borel functions is equivalent to the transfinite leftmost path principle, which falls strictly between ATR_0 and $\Pi_1^1\text{-CA}_0$. We also exhibit several weakenings of Caristi's theorem that are equivalent to WKL_0 and to ACA_0 .

1. INTRODUCTION

Metric fixed point theorems state that, under certain conditions, a function $f: \mathcal{X} \rightarrow \mathcal{X}$ from a metric space to itself has a fixed point, i.e. there is an $x_* \in \mathcal{X}$ such that $f(x_*) = x_*$. Such theorems have many applications in geometry, partial differential equations, etc., where the function f is typically continuous.

One fixed point theorem that does *not* require the continuity of f is Caristi's fixed point theorem [1]. Instead, the function f is 'controlled' by a non-negative lower semi-continuous function. Specifically, it applies to what we will call 'Caristi systems'. As is typical, we will notationally identify a metric space (\mathcal{X}, d) with \mathcal{X} . Recall that $V: \mathcal{X} \rightarrow \mathbb{R}$ is lower semi-continuous if whenever $x_n \rightarrow x$, it follows that $V(x) \leq \liminf_{n \rightarrow \infty} V(x_n)$.

Definition 1.1. A *Caristi system* is a tuple (\mathcal{X}, f, V) , where \mathcal{X} is a complete metric space, $f: \mathcal{X} \rightarrow \mathcal{X}$ is arbitrary, $V: \mathcal{X} \rightarrow [0, \infty)$ is lower semi-continuous, and for all $x \in \mathcal{X}$, $d(x, f(x)) \leq V(x) - V(f(x))$.

Theorem 1.2 (Caristi [1]). *Every Caristi system (\mathcal{X}, f) has a fixed point; i.e. there is $x_* \in \mathcal{X}$ such that $f(x_*) = x_*$.*

Caristi's theorem has various applications and generalizations in metric fixed point theory [6, 7]. We think of $V: \mathcal{X} \rightarrow [0, \infty)$ as a 'potential', which diminishes after applying f ; intuitively, after enough applications of f , no more potential is lost and we have reached a fixed point. In Caristi's original proof, f is iterated transfinitely, but this can be avoided by using Ekeland's variational principle.

Theorem 1.3 (Ekeland [3]). *Let \mathcal{X} be a complete metric space and $V: \mathcal{X} \rightarrow [0, \infty)$ be lower semi-continuous. Then there is an $x_* \in \mathcal{X}$ such that for all $x \in \mathcal{X}$, $d(x_*, x) \leq V(x_*) - V(x)$ implies that $x = x_*$.*

We call such an x_* a *critical point* of V . Theorem 1.2 can be derived from Theorem 1.3 by observing that any critical point for V will also be a fixed point of f , by the assumption that $d(x_*, f(x_*)) \leq V(x_*) - V(f(x_*))$. Note however that Ekeland's theorem is equivalent to $\Pi_1^1\text{-CA}_0$, so the transfinite methods are still hidden 'under the hood' [4].

Another fixed point theorem that was originally proven via infinitary methods (stated in terms of an explicit invocation of Zorn's lemma) is the Priess-Crampe & Ribenboim theorem, which deals with spherically complete ultrametric spaces. This result is partially motivated by logic programming [10] and has recently found applications in cut-elimination for ill-founded proofs [12].

Definition 1.4. A metric space \mathcal{X} is an *ultrametric space* if for all $x, y, z \in \mathcal{X}$, $d(x, y) \leq \max\{d(x, z), d(z, y)\}$. \mathcal{X} is *spherically complete* if whenever $\langle \overline{B}_{\rho_i}(x_i) \rangle_{i \in \mathbb{N}}$ is a decreasing sequence of closed balls, it follows that $\bigcap_{i \in \mathbb{N}} \overline{B}_{\rho_i}(x_i) \neq \emptyset$.

A function $f: \mathcal{X} \rightarrow \mathcal{X}$ is *strictly contracting* if for all $x \neq y \in \mathcal{X}$, $d(f(x), f(y)) < d(x, y)$.

Theorem 1.5 (Priess-Crampe & Ribenboim [11]). *Let \mathcal{X} be a spherically complete ultrametric space and $f: \mathcal{X} \rightarrow \mathcal{X}$ be strictly contracting. Then, f has a unique fixed point.*

Key words and phrases. computability theory, reverse mathematics, second-order arithmetic, fixed-point theorems, variational principles.

Our goal is to determine the strength of the Priess-Crampe & Ribenboim theorem and Caristi's theorem in the sense of reverse mathematics.

For Caristi's theorem, in order to deal with 'arbitrary' f , we consider the case where f is either Baire or Borel, as these are wide classes that can readily be coded within second-order arithmetic. The Caristi theorem for these classes is strictly between ATR_0 and $\Pi_1^1\text{-CA}_0$, and indeed equivalent to the theory of the transfinite leftmost path principle introduced by Towsner [14]. This shows that transfinite methods cannot be avoided altogether. The reversal, however, requires a fairly complicated choice of \mathcal{X} and f : Caristi's theorem only requires such a strong theory because it covers such complicated functions on fairly general spaces.

If we restrict \mathcal{X} or f to nicer examples, Caristi's theorem becomes easier to prove. The case where both f and V are continuous has also been treated by Peng and Yamazaki [9]; this case is interesting, as it can already be viewed as a generalization of the Banach fixed point theorem [6]. Using our previous work on the reverse mathematics of Ekeland's variational principle [4], we extend this treatment to lower semi-continuous V . As we will see, weakened versions of Caristi's theorem are equivalent to either WKL_0 when \mathcal{X} is compact and both functions are continuous, or ACA_0 when either compactness [9] or continuity is dropped (but not both).

Regarding the Priess-Crampe & Ribenboim theorem, surprisingly it may already be proven in RCA_0 . We show this by exhibiting a new constructive proof.

2. SUBSYSTEMS OF SECOND-ORDER ARITHMETIC

We will work within subsystems of second-order arithmetic as in [13]. The language is that of Peano arithmetic enriched with variables for sets of natural numbers which may be quantified over. We use Δ_0^0 to denote the set of all formulas, possibly with set parameters, where no second-order quantifiers appear and all first-order quantifiers are bounded, and as usual define the classes Σ_n^e and Π_n^e where n is the number of alternating first-order (for $e = 0$) or second-order (for $e = 1$) quantifiers (see e.g. [13] for details).

We use the notation $\langle x_0, \dots, x_n \rangle$ to denote sequences of natural numbers encoded in a standard way. As usual, sets of pairs may be used to represent binary relations and functions on the natural numbers, and for a binary relation R , $|R|$ denotes the union of the domain and codomain of R .¹ The set of all finite sequences of natural numbers is denoted $\mathbb{N}^{<\mathbb{N}}$. For $\sigma, \tau \in \mathbb{N}^{<\mathbb{N}}$ we write $\sigma \sqsubseteq \tau$ if σ is an initial segment of τ , $\sigma \sqsubset \tau$ if σ is a proper initial segment of τ , and set $\downarrow \sigma = \{\tau \in \mathbb{N}^{<\mathbb{N}} : \tau \sqsubseteq \sigma\}$. If $\Lambda : \mathbb{N} \rightarrow \mathbb{N}$ and $n \in \mathbb{N}$, write $\Lambda \upharpoonright n$ for the finite sequence $\langle \Lambda(i) \rangle_{i < n}$; note that the sequence is empty when $n = 0$. For a set $X \subseteq \mathbb{N}$ we write $X \upharpoonright n$ for the finite sequence $\lambda_X \upharpoonright n$ where λ_X is the characteristic function of X . We extend the use of \sqsubseteq by defining $\sigma \sqsubseteq \Lambda$ whenever $\sigma = \Lambda \upharpoonright n$ for some n . Concatenation of sequences is denoted by \frown . If $X, Y \subseteq \mathbb{N}$ then $X \oplus Y$ is $\{2n : n \in X\} \cup \{2n + 1 : n \in Y\}$, the Turing join of X and Y .

We will represent trees as subsets of $\mathbb{N}^{<\mathbb{N}}$ which are downward closed under \sqsubseteq . Binary trees are then those trees which are subsets of $\{0, 1\}^{<\mathbb{N}}$. An infinite sequence Λ is a *path through* T if for every n , $\Lambda \upharpoonright n \in T$, and $[T]$ denotes the class of paths through T .

If \prec is a well-order on a subset of \mathbb{N} , then $\text{ot}(\prec)$ denotes the order-type of \prec . If moreover $X \subseteq \mathbb{N}^2$ is a set of pairs, and $\alpha \in |\prec|$, we write X_α for $\{x \in \mathbb{N} : \langle x, \alpha \rangle \in X\}$ and $X_{\prec \alpha}$ for the set of all $\langle n, \beta \rangle \in X$ with $\beta \prec \alpha$, with $X_{\preceq \alpha}$ being defined analogously. Transfinite recursion along \prec may be defined in second-order arithmetic, as follows.

Definition 2.1 ([13, Chapter V]). Let $\theta(x, Y, \vec{z}, \vec{Z})$ be any formula. We define $H_\theta(\prec, Y, \vec{z}, \vec{Z})$ to be the formula which says that, for each $\alpha \in |\prec|$, $Y_\alpha = \{x : \theta(x, Y_{\prec \alpha}, \vec{z}, \vec{Z})\}$ and for $\alpha \notin |\prec|$, $Y_\alpha = \emptyset$. We may write $H_\theta(\prec, Y)$ when \vec{z}, \vec{Z} are clear from context.

We may choose θ so that for any parameters Y, \vec{z}, \vec{Z} , the set $\{x : \theta(x, Y, \vec{z}, \vec{Z})\}$ is a universal computably enumerable set relative to Y, \vec{Z} —that is, so that for any Y, \vec{z}, \vec{Z} , any set computably enumerable relative to Y, \vec{Z} is equal to some slice $\{x : \theta(\langle e, x \rangle, Y, \vec{z}, \vec{Z})\}$. In this case the choice of θ only matters up to details of coding, so we omit it.

Definition 2.2. For a fixed θ_* such that for any parameters Y, \vec{z}, \vec{Z} , the set $\{x : \theta_*(x, Y, \vec{z}, \vec{Z})\}$ is a universal computably enumerable set relative to Y, \vec{Z} , we write H instead of H_{θ_*} .

¹Within RCA_0 , $|R|$ may not exist as a set but one can always find an isomorphic relation R' such that $|R'|$ exists as a set.

This suffices to define the ‘Big Five’ theories of reverse mathematics: RCA_0 includes basic axioms of arithmetic together with induction for Σ_1^0 -definable predicates and comprehension for Δ_1^0 -definable predicates; WKL_0 extends RCA_0 with the formalized weak König’s lemma; and ACA_0 includes comprehension for arithmetical formulas. Then ATR_0 ensures that, whenever \prec is a well-order, there is a unique Y so that $H(\prec, Y, \vec{z}, \vec{Z})$ holds, and finally, $\Pi_1^1\text{-CA}_0$ is axiomatized with comprehension for Π_1^1 formulas. We have mentioned these theories in strictly increasing order of strength, but there is a less known theory between ATR_0 and $\Pi_1^1\text{-CA}_0$ due to the third author [14].

Definition 2.3. When \prec is a linear order, we say \prec is a *successor* if it has a maximal element x , and we define its *predecessor* $\prec^- = \prec \upharpoonright (|\prec| \setminus \{x\})$. When \prec is a well-order, we say W is Σ_2^Z if either (a) \prec is a successor and W is computably enumerable in the unique $\bar{Y} := Y \oplus Z$ so that $H(\prec^-, Y, Z)$ holds, or (b) \prec is not a successor and W is computable in the $\bar{Y} := Y \oplus Z$ so that $H(\prec, Y, Z)$ holds.²

TLPP_0 is defined to be RCA_0 together with the TLPP (transfinite leftmost path principle) axiom. When Λ and Γ are infinite sequences, we write $\Gamma < \Lambda$ if there is some n such that $\Gamma \upharpoonright n = \Lambda \upharpoonright n$ and $\Gamma(n) < \Lambda(n)$. Then TLPP is the formalization of the following statement: *whenever $T \subseteq \mathbb{N}^{<\mathbb{N}}$ is a tree with an infinite path and \prec is a well-order, there is a path Λ through T such that there is no path Γ through T which is $\Sigma_2^{\Lambda \oplus T}$ and $\Gamma < \Lambda$.* We call Λ a *relativized \prec -leftmost path* for T , or just *relativized leftmost path* when \prec, T are clear from context. For a detailed treatment of these and other subsystems of second-order arithmetic, see [13, 14]. The following two characterizations of TLPP_0 will be useful; the proof is simply a relativization of the one in [8].

Lemma 2.4 (RCA_0). *The following are equivalent:*

- (1) TLPP_0 .
- (2) *Whenever $\langle T_n \rangle_{n \in \mathbb{N}}$ is a sequence of trees and \prec is a well-order, there are a pair of sets Z_0, Z_1 so that $n \in Z_0$ if and only if there is a path through T_n which is $\Sigma_2^{Z_0 \oplus Z_1 \oplus \langle T_n \rangle_{n \in \mathbb{N}}}$.*
- (3) *For any parameters \vec{Z} , any well-order \prec , and any Σ_2^0 formula ϕ , there are sets X_0, X_1 so that $x \in X_0$ iff for every Y which is $\Sigma_2^{X_0 \oplus X_1 \oplus \vec{Z}}$, $\phi(x, Y, \vec{Z})$ holds.*

Proof. To show 1 implies 2, let $\langle T_n \rangle_{n \in \mathbb{N}}$ be given. We define a single tree T intertwining these trees: we first define $T'_n = \{\langle 0 \rangle \frown \sigma : \sigma \in T_n\} \cup \{\sigma : \forall i < |\sigma| \sigma(i) = 1\}$ (that is, we add a single infinite path to the right of all paths in T_n) and then take $T = \{\sigma : \forall \langle i, n \rangle < |\sigma| \langle \sigma(\langle 0, n \rangle), \dots, \sigma(\langle i, n \rangle) \rangle \in T'_n\}$. By TLPP_0 , let Λ be a relativized leftmost path for T' and let $Z = \{n : \Lambda(\langle 0, n \rangle) = 0\}$.

If $n \in Z$ then there is a path through T_n computable from Λ . If $n \notin Z$ and Γ is a path through T_n which is $\Sigma_2^{Z \oplus \Lambda \oplus \langle T_n \rangle_{n \in \mathbb{N}}}$, then we can modify Λ by setting $\Lambda'(\langle 0, n \rangle) = 0$, $\Lambda'(\langle i+1, n \rangle) = \Gamma(i)$, and $\Lambda'(\langle i, m \rangle) = \Lambda(\langle i, m \rangle)$ for $m \neq n$. But $\Lambda' < \Lambda$ and Λ' is $\Sigma_2^{\Lambda \oplus \langle T_n \rangle_{n \in \mathbb{N}}}$, contradicting the choice of Λ .

Next we show that 2 implies 3. By the normal form theorem, write $\phi(x, Y, \vec{Z}) \equiv \neg \forall w \exists z \psi(w, z, x, Y \upharpoonright z, \vec{Z})$. For each x , let T_x be the tree defined by $\sigma \in T_x$ iff for all $i \leq j < |\sigma|$, $\sigma(i) \sqsubseteq \sigma(j) \in 2^{\mathbb{N}}$, $i \leq |\sigma(i)|$ and $\exists z \leq |\sigma(i)| \psi(i, z, x, \sigma(i) \upharpoonright z, \vec{Z})$ holds.

We extend $\langle T_x \rangle_{x \in \mathbb{N}}$ to encode \vec{Z} and find sets S_0, S_1 so that $x \in S_0$ iff there is a path through T_x which is $\Sigma_2^{S_0 \oplus S_1 \oplus \langle T_x \rangle_{x \in \mathbb{N}} \oplus \vec{Z}}$. The complement \bar{S}_0 of S_0 is the desired set. If $x \notin \bar{S}_0$, then there is a path Λ through T_x which is $\Sigma_2^{S_0 \oplus S_1 \oplus \langle T_x \rangle_{x \in \mathbb{N}} \oplus \vec{Z}}$. Since $\langle T_x \rangle_{x \in \mathbb{N}}$ is computable from \vec{Z} , this path is $\Sigma_2^{\bar{S}_0 \oplus S_1 \oplus \vec{Z}}$ and $Y = \{i : \Lambda(i)(i) = 1\}$ witnesses $\forall w \exists z \psi(w, z, x, Y \upharpoonright z, \vec{Z})$. On the other hand, if there is a Y so that $\forall w \exists z \psi(w, z, x, Y \upharpoonright z, \vec{Z})$ and Y is $\Sigma_2^{\bar{S}_0 \oplus S_1 \oplus \vec{Z}}$ then a function Λ defined by $\Lambda(i) = Y \upharpoonright z_i$ where z_i is the smallest $z > i$ such that $\psi(i, z, x, Y \upharpoonright z, \vec{Z})$ holds is a path through T_x which is $\Sigma_2^{S_0 \oplus S_1 \oplus \langle T_x \rangle_{x \in \mathbb{N}} \oplus \vec{Z}}$ and therefore $x \notin \bar{S}_0$.

To show that 3 implies 1, let T be a tree with an infinite path P , and let \prec be a well-order. Let $\phi(\sigma, Y, T)$ be a formula which holds iff Y is not a path through $T_\sigma = \{\tau \in T : \sigma \sqsubseteq \tau\}$. By 3, let X_0 and X_1 be such that $\sigma \in X_0$ iff no Y which is $\Sigma_2^{X_0 \oplus X_1 \oplus T \oplus P}$ is a path through T extending σ . Now recursively compute a path Λ from X_0 by setting $\Lambda(n)$ to be the least i such that $(\Lambda \upharpoonright n) \frown \langle i \rangle \notin X_0$. The value $\Lambda(0)$ is defined because $P \upharpoonright 1 \notin X_0$. If $\Lambda \upharpoonright n$ is defined for some $n \geq 1$, then $\Lambda \upharpoonright n \notin X_0$, so there is a $\Sigma_2^{X_0 \oplus X_1 \oplus T \oplus P}$ path Y through T extending $\Lambda \upharpoonright n$. Then $Y \upharpoonright (n+1) \notin X_0$, which implies that $\Lambda(n)$ is defined and therefore that $\Lambda \upharpoonright (n+1)$ is defined. Thus Λ is a path through T . Now suppose for

²Note that there is an irregularity at the lowest level in our notation, namely, W is Σ_0^Z means that W is Δ_1^Z and not Σ_0^Z in the sense that W is defined by a bounded formula from Z .

a contradiction that there is a $\Sigma_{\prec}^{\Lambda \oplus T}$ path Γ through T with $\Gamma < \Lambda$. Let n be such that $\Gamma \upharpoonright n = \Lambda \upharpoonright n$ and $\Gamma(n) < \Lambda(n)$. Then $\Gamma \upharpoonright (n+1) \notin X_0$ because Γ is $\Sigma_{\prec}^{\Lambda \oplus T}$ and hence $\Sigma_{\prec}^{X_0 \oplus X_1 \oplus T \oplus P}$. This contradicts the definition of $\Lambda(n)$ because $(\Lambda \upharpoonright n) \frown \langle \Gamma(n) \rangle = \Gamma \upharpoonright (n+1) \notin X_0$, but $\Gamma(n) < \Lambda(n)$. \square

The unrelativized version of (iii)—the existence of a set X so that $x \in X$ iff for every Y , $\phi(x, Y, \vec{Z})$ holds, is sometimes called an *impredicative* definition (since the set Y might be X itself, or something defined from X). The relativized version is called a *partial impredicativity* because we only consider those Y 's which are defined from X in a limited way.

3. BAIRE AND BOREL FUNCTIONS

Part of the appeal of second-order arithmetic as a foundational system for mathematics is that it suffices to develop a large part of mathematical analysis, particularly when dealing with separable metric spaces. However, this requires some coding machinery. In this section, we recall this machinery, and establish notation that will be used throughout the text.

Definition 3.1 (RCA₀; see [13, Definition II.5.1]). A (code for a) complete separable metric space $\mathcal{X} = \widehat{X}$ is defined in RCA₀ to be a nonempty set $X \subseteq \mathbb{N}$ together with a sequence of real numbers $d: X \times X \rightarrow \mathbb{R}$ such that $d(a, a) = 0$, $d(a, b) = d(b, a) \geq 0$, and $d(a, b) + d(b, c) \geq d(a, c)$ for all $a, b, c \in X$. A *point* of \widehat{X} is a sequence $x = \langle x_i \rangle_{i \in \mathbb{N}}$ of elements of X such that for all $i \leq j$, $d(x_i, x_j) \leq 2^{-i}$. We write $x \in \widehat{X}$ to mean that x is a point of \widehat{X} . We set $d(x, y) = \lim_{n \rightarrow \infty} d(x_n, y_n)$, which provably exists in RCA₀.

We say that the space \mathcal{X} is *compact* if there is a sequence of points witnessing that \mathcal{X} is totally bounded [13, Definition III.2.3]. For our purposes, it suffices to mention that RCA₀ proves that both $[0, 1]$ and the Cantor space are compact in this sense. Here, the Cantor space is $\{0, 1\}^{\mathbb{N}}$ with $d(\Lambda, \Lambda') = 2^{-n}$, for the least n such that $\Lambda(n) \neq \Lambda'(n)$ and $d(\Lambda, \Lambda') = 0$ when no such n exists. The *Baire space* is defined analogously, but with set of points $\mathbb{N}^{\mathbb{N}}$; note that the Baire space is *not* compact. In the definitions below, $\mathbb{Q}^{>0}$ denotes $\{q \in \mathbb{Q} : q > 0\}$.

Definition 3.2 (RCA₀; see [13, Definition II.5.1]). Let \widehat{X} be a complete separable metric space. The (code for the) *rational open ball* $B_r(a)$ is the ordered pair $\langle a, r \rangle$, with $a \in X$ and $r \in \mathbb{Q}^{>0}$. We define $B_r(a) \Subset B_q(b)$ if $d(a, b) + r < q$ and $B_r(a) \subseteq B_q(b)$ if $d(a, b) + r \leq q$.

We remark that $B_r(a) \Subset B_q(b)$ implies that $B_r(a) \subseteq B_q(b)$ (in the usual set-theoretic sense), but not necessarily the converse (consider e.g. the case where \mathcal{X} is a singleton).

One challenge in formalizing Caristi's theorem in the context of second-order arithmetic is that it applies to *arbitrary* functions f , which would in principle require a third-order quantification. Instead, we will work with a rather wide class of functions that can still be formalized as second-order objects: Baire and Borel functions. To keep our presentation unified and to minimize coding concerns, we view all functions on metric spaces as special cases of Baire or Borel functions, which is not how e.g. continuous or lower semi-continuous functions are typically coded in the literature. In Appendix A we discuss the relationship between our codes and the more standard ones.

The general theory of Borel sets in reverse mathematics is well established, for instance in [13, 15], but Borel *functions* have been less well studied.

Definition 3.3. When \mathcal{X} is a complete separable metric space, recall that a *Borel code* (that is, a code for a Borel subset of \mathcal{X}) is a tree of sequences S such that: (a) there is no infinite path through S , (b) there is exactly one n so that $\langle n \rangle \in S$, and (c) any leaf $\sigma \in S$ has the form $\tau \frown \langle B_r(a) \rangle$, where $a \in X$ and $r \in \mathbb{Q}^{>0}$.

We view such a tree as coding a Borel set. In fact, every node in the tree will code a Borel set $U_S(\sigma)$ defined recursively by

- when $\sigma = v \frown \langle B_r(a) \rangle$ is a leaf, $U_S(\sigma) = B_r(a)$,
- when $\sigma = v \frown \langle n \rangle$ where n is odd, $U_S(\sigma) = \bigcup \{U_S(\sigma \frown \langle k \rangle) : \sigma \frown \langle k \rangle \in S\}$,
- when $\sigma = v \frown \langle n \rangle$ where n is even, $U_S(\sigma) = \bigcap \{U_S(\sigma \frown \langle k \rangle) : \sigma \frown \langle k \rangle \in S\}$,
- when $\sigma = \langle \rangle$, $U_S(\sigma) = U_S(\langle n \rangle)$, where n is unique so that $\langle n \rangle \in S$.

In particular, open sets may be represented as unions of open balls, hence via a Borel code. By abuse of notation, we write $x \in S$ as an abbreviation for $x \in U_S(\langle \rangle)$. Note that the latter is not a

formula of second-order arithmetic, however it is a theorem that, in ATR_0 , we can identify membership in coded Borel sets (see [13, Lemma V.3.3 and Definition V.3.4]).

Lemma 3.4 (ATR_0). *If $\langle x_n \rangle_{n \in \mathbb{N}}$ is a sequence of points in \mathcal{X} and $\langle S_n \rangle_{n \in \mathbb{N}}$ is a sequence of Borel codes then $\{n : x_n \in U_{S_n}(\langle \rangle)\}$ exists.*

We will need to discuss Borel functions. We encode a Borel function as one where the inverse image of each basic open set is given by a Borel code.

Definition 3.5. Let \mathcal{X} and \mathcal{Y} be metric spaces and Υ be a set coding a sequence of Borel codes $\langle \Upsilon_{\langle a,r \rangle} : \langle a,r \rangle \in Y \times \mathbb{Q}_{>0} \rangle$. Write $\Upsilon(B_r(a))$ for $U_{\Upsilon_{\langle a,r \rangle}}(\langle \rangle)$.

We say that Υ is a (code for) a Borel function from \mathcal{X} to \mathcal{Y} if for all basic open balls B, B' of \mathcal{Y} :

- (i) if $B \subseteq B'$ then $\Upsilon(B) \subseteq \Upsilon(B')$,
- (ii) if $B \cap B' = \emptyset$ then $\Upsilon(B) \cap \Upsilon(B') = \emptyset$,

The set Υ codes the partial function $f : \mathcal{X} \rightarrow \mathcal{Y}$, where $x \in \text{dom}(f)$ if there is a (unique) element $f(x) = y \in \mathcal{Y}$ such that for every ball B containing y , $x \in \Upsilon(B)$.

Recall that the Kleene-Brouwer order of a tree T , denoted $\text{KB}(T)$, is a linearization of T which, provably in ACA_0 , remains well founded when T is (see e.g. [13]). When f is a Borel function coded by Υ , the complexity of f , written $\|f\|$, is $\text{KB}(\{\langle n \rangle \frown \sigma : \sigma \in \Upsilon_n\})$.

In particular, continuous functions may be coded as Borel functions, where the preimage of every open ball is open.

Lemma 3.6 (ATR_0). *Let Υ be a code for a Borel function from \mathcal{X} to \mathcal{Y} . Then, for any $x \in \mathcal{X}$, the image of x under f exists and is $\Sigma_{\|f\|}^{x \oplus \Upsilon}$.*

Proof. The evaluation function of the set of balls $B_r(a)$ such that $x \in f^{-1}(B_r(a))$ is $\Sigma_{\|f\|}^{x \oplus \Upsilon}$. Therefore we may define an approximation by taking y_i to be the least a such that $x \in f^{-1}(B_{2^{-i-1}}(a))$, and then the sequence $\langle y_i \rangle_{i \in \mathbb{N}}$ converges to $f(x)$. \square

Rather than referring to codes, we will usually talk about the function f : saying that f is coded is simply saying that, for each $B_r(a)$, the set $f^{-1}(B_r(a))$ is given by a Borel code, and these codes are presented uniformly in a, r .

Definition 3.7. A (code for a) partial Baire function is a well-founded tree Ξ such that each leaf is labeled by a (code for a) partial continuous function and each non-leaf has an extension for each $i \in \mathbb{N}$.

For any x and any $\sigma \in \Xi$, the value $f_{\Xi, \sigma}(x)$, if it exists, is defined recursively by:

- if σ is a leaf labeled by f then $f_{\Xi, \sigma}(x) = f(x)$,
- if σ is not a leaf and there is any n so that $f_{\Xi, \sigma \frown \langle n \rangle}(x)$ does not exist then $f_{\Xi, \sigma}(x)$ does not exist,
- if σ is not a leaf, $f_{\Xi, \sigma \frown \langle n \rangle}(x)$ exists for all n , then $f_{\Xi, \sigma}(x) = \lim_{n \rightarrow \infty} f_{\Xi, \sigma \frown \langle n \rangle}(x)$ if this exists, and does not exist otherwise.

We write $f_{\Xi}(x)$ for $f_{\Xi, \langle \rangle}(x)$ and call f a Baire function if it is given by a code for a partial Baire function such that, for every x , $f(x)$ exists.

It is provable in ATR_0 that every Baire function is Borel, and the converse is true on zero-dimensional spaces, including the Cantor space and Baire space; see Appendix A for details.

Potentials (i.e., non-negative lower semi-continuous functions) can be represented as increasing limits of continuous functions and hence as Baire class 1 functions. For this, we use the following lemma (not formalized in second-order arithmetic).

Lemma 3.8. *If \mathcal{X} is any metric space and V is a potential on \mathcal{X} , then there exists a pointwise increasing sequence of continuous functions $V_n : \mathcal{X} \rightarrow [0, \infty)$ such that $V = \lim_{n \rightarrow \infty} V_n$.*

Proof sketch. It is easy to check that an increasing limit of continuous functions is lsc. For the converse, given $\alpha \geq 0$, define the α -envelope of V , denoted $V_{(\alpha)}$, by $V_{(\alpha)}(x) = \inf_{y \in \mathcal{X}} (V(y) + \alpha d(x, y))$. Then, it is not hard to check that $(V_{(n)})_{n \in \mathbb{N}}$ is an increasing sequence of continuous functions converging pointwise to V . \square

Thus in second-order arithmetic we define a potential to be a Baire class 1 function which is an limit of pointwise-monotone continuous functions V_n (i.e. $V_n(x) \leq V_m(x)$ if $n \leq m$). If V_n is the n -envelope

of V , we say that V is *enveloped*. Note that $V_{(\alpha)}$ is well defined even when $\alpha = 0$, in which case we obtain $V_{(0)} = \inf V$, and thus enveloped potentials have an infimum. As we will see, there is much more information that can be extracted from continuous envelopes.

This coding machinery will suffice to formalize Caristi's theorem, but first we turn our attention to Priess-Crampe, which involves only continuous functions.

4. AN ELEMENTARY PROOF OF THE PRIESS-CRAMPE & RIBENBOIM THEOREM

Recall that Theorem 1.5 states that if \mathcal{X} is a spherically complete ultrametric space and $f: \mathcal{X} \rightarrow \mathcal{X}$ is strictly contracting, then f has a unique fixed point. This theorem may be stated in second-order arithmetic via the coding machinery described above. For spherical completeness, let $\overline{B}_\rho(x)$ denote the class of $y \in \mathcal{X}$ such that $d(x, y) \leq \rho$ for given $x \in \mathcal{X}$ and $\rho > 0$. As \mathcal{X} is an ultrametric space, we write $\overline{B}_\rho(x) \subseteq \overline{B}_\delta(y)$ to denote that $\max\{d(x, y), \rho\} \leq \delta$ and observe that $\overline{B}_\rho(x) \subseteq \overline{B}_\delta(y)$ implies that $\overline{B}_\rho(x) \subseteq \overline{B}_\delta(y)$. Then, \mathcal{X} is spherically complete if whenever $\langle \overline{B}_{\rho_i}(x_i) \rangle_{i \in \mathbb{N}}$ is a sequence of closed balls such that $\overline{B}_{\rho_i}(x_i) \supseteq \overline{B}_{\rho_{i+1}}(x_{i+1})$ for all i , there is an $x \in \bigcap_{i \in \mathbb{N}} \overline{B}_{\rho_i}(x_i)$. We now obtain the following.

Theorem 4.1. *The Priess-Crampe & Ribenboim theorem is provable in RCA_0 .*

Proof. Uniqueness follows easily from the assumption that f is strictly contracting, so we focus on existence.

Let $\mathcal{X} = \widehat{X}$, and let $\langle a_i \rangle_{i \in \mathbb{N}}$ enumerate X with each a_i occurring infinitely often. Define $\rho(x) = d(x, f(x))$. We claim that there exists a sequence $\langle b_i \rangle_{i \in \mathbb{N}}$ such that $\rho(b_i) \geq \rho(b_{i+1})$ for all i and $b_i \rightarrow \inf \rho$ as $i \rightarrow \infty$ (in the sense that for all $x \in \mathcal{X}$ and $\varepsilon > 0$, there is an i with $\rho(b_i) < \rho(x) + \varepsilon$). Construct the sequence $\langle b_i \rangle_{i \in \mathbb{N}}$ as follows. Note that $\rho(x) < \rho(y)$ is a Σ_1^0 statement, which we may represent as $\exists z \phi(x, y, z)$. Let $b_0 = a_0$, and recursively define $b_{i+1} = a_{i+1}$ if there is a $z < i$ witnessing that $\rho(a_{i+1}) < \rho(b_i)$ (in the sense that $\phi(a_{i+1}, b_i, z)$ holds), and otherwise define $b_{i+1} = b_i$. It is not hard to check that the sequence $\langle b_i \rangle_{i \in \mathbb{N}}$ satisfies the required properties.

Let $\rho_i = \rho(b_i)$ for each i . We show that $\overline{B}_{\rho_i}(b_i) \supseteq \overline{B}_{\rho_{i+1}}(b_{i+1})$ for all i . To see this, observe that

$$\begin{aligned} d(b_i, b_{i+1}) &\leq \max\{d(b_i, f(b_i)), d(f(b_i), f(b_{i+1})), d(b_{i+1}, f(b_{i+1}))\} \\ &= \max\{\rho_i, d(f(b_i), f(b_{i+1})), \rho_{i+1}\} = \rho_i. \end{aligned}$$

The last equality holds because f is strictly contracting and therefore $d(f(b_i), f(b_{i+1})) < d(b_i, b_{i+1})$. Thus it must be that $\max\{\rho_i, d(f(b_i), f(b_{i+1})), \rho_{i+1}\} = \max\{\rho_i, \rho_{i+1}\} = \rho_i$. It follows that $\max\{d(b_i, b_{i+1}), \rho_{i+1}\} \leq \rho_i$, so $\overline{B}_{\rho_i}(b_i) \supseteq \overline{B}_{\rho_{i+1}}(b_{i+1})$.

By spherical completeness, there is an $x_* \in \bigcap_{i \in \mathbb{N}} \overline{B}_{\rho_i}(b_i)$. Then $\rho(x_*) \leq \rho_i$ for all i because

$$d(x_*, f(x_*)) \leq \max\{d(x_*, b_i), d(b_i, f(b_i)), d(f(b_i), f(x_*))\} = \max\{d(x_*, b_i), \rho_i\} = \rho_i,$$

since $x \in \overline{B}_{\rho_i}(b_i)$ and f is strictly contracting. If $\rho(x_*) > 0$ then $d(x_*, f(x_*)) > d(f(x_*), f^2(x_*)) = \rho(f(x_*))$, which contradicts the fact that $\lim_{i \rightarrow \infty} \rho(b_i) = \inf \rho$. We conclude that $d(x_*, f(x_*)) = 0$. \square

5. THE CARISTI THEOREM IN THE BIG FIVE

In this section we study weakenings of Caristi's theorem provable in WKL_0 and ACA_0 . Caristi's theorem follows directly from Ekeland's variational principle, so we first recall the main results from [4] regarding the reverse mathematics of the latter. In particular, Ekeland's variational principle is derivable in $\Pi_1^1\text{-CA}_0$, thus establishing an upper bound for Caristi's theorem. Ekeland's variational principle also has natural weakenings derivable in WKL_0 and in ACA_0 .

Definition 5.1. Given definable classes \mathfrak{X} of coded metric spaces and \mathfrak{V} of coded potentials, the (formalized) *Ekeland variational principle* for $\mathcal{X} \in \mathfrak{X}$ and $V \in \mathfrak{V}$ is the statement that, if $\mathcal{X} \in \mathfrak{X}$ is a coded separable complete metric space and $V \in \mathfrak{V}$ is a coded potential, then there exists $x_* \in \mathcal{X}$ such that for all $x \in \mathcal{X}$, if $d(x_*, x) \leq V(x_*) - V(x)$, then $x = x_*$.

When not mentioned, we assume that \mathfrak{X} is the class of all coded complete separable metric spaces and \mathfrak{V} is the class of all coded potentials.

The following is proven in [4], and in fact all items reverse.

Theorem 5.2. *The Ekeland variational principle holds:*

- (i) (WKL_0) For compact \mathcal{X} and continuous V .
- (ii) (ACA_0) For compact \mathcal{X} or continuous V .

(iii) (Π_1^1 -CA₀) For the class of all metric spaces with an arbitrary potential.

All of these results reverse, although we won't be needing this. We remark that [4] uses a different presentation of potentials (i.e. lower semi-continuous functions), but the two are equivalent over RCA₀ (see Appendix A). The discontinuous cases are established by reducing to continuous cases via envelopes, given the following [4, Lemma 8.1].

Lemma 5.3 (RCA₀). *Let V be any potential and $\alpha > 1$ be such that $V_{(\alpha)}$ exists. Then, any critical point x_* of $V_{(\alpha)}$ is also a critical point of V , and $V_{(\alpha)}(x_*) = V(x_*)$.*

We are now ready to state our formalization of the Caristi fixed point theorem and prove it (and its weakenings) in standard systems of second-order arithmetic.

Definition 5.4. A *Caristi system* is a tuple (\mathcal{X}, f, V) , where \mathcal{X} is a coded complete separable metric space, $f: \mathcal{X} \rightarrow \mathcal{X}$ a coded Baire or Borel function, and $V: \mathcal{X} \rightarrow [0, \infty)$ a coded potential.

Given definable classes \mathfrak{X} of complete, separable metric spaces and $\mathfrak{F}, \mathfrak{V}$ of functions such that elements of \mathfrak{F} are Borel or Baire functions of the form $f: \mathcal{X} \rightarrow \mathcal{X}$ and elements of \mathfrak{V} are potentials of the form $V: \mathcal{X} \rightarrow [0, \infty)$, the (*formalized*) *Caristi fixed point theorem (CFP)* for $\mathcal{X} \in \mathfrak{X}$, $f \in \mathfrak{F}$, and $V \in \mathfrak{V}$ is the statement that, if (\mathcal{X}, f, V) is a Caristi system whose elements belong to the aforementioned classes, then there exists $x_* \in \mathcal{X}$ such that $x_* = f(x_*)$.

When not mentioned, we assume that \mathfrak{X} is the class of all coded separable complete metric spaces, \mathfrak{F} is the class of all Baire or Borel functions, and \mathfrak{V} is the class of all coded potentials.

The following is a corollary of Theorem 5.2, using the fact that the Ekeland variational principle implies Caristi's theorem.

Proposition 5.5. *The CFP holds whenever:*

- (i) (WKL₀) \mathcal{X} is compact and V is continuous.
- (ii) (ACA₀) \mathcal{X} is compact, f is continuous, or V is continuous.
- (iii) (Π_1^1 -CA₀) Always.

Proof. Most items follow by using Theorem 5.2 and the fact that every critical point of V is a fixed point of f . The exception is (ii) for continuous f , but the proof of Peng and Yamazaki [9], itself a version of the classic proof of the Banach fixed point theorem, works in this context.

Let (\mathcal{X}, f, V) be a Caristi system where f is continuous and $x_0 \in \mathcal{X}$. Define a sequence x_n given recursively by $x_{n+1} = f(x_n)$; this sequence is readily available in ACA₀ using the continuity of f . Similarly, the sequence given by $v_n = V(x_n)$ exists, as ACA₀ can compute suprema uniformly. Since $d(x_n, f(x_n)) \leq V(x_n) - V(f(x_n)) = V(x_n) - V(x_{n+1})$, we must have that $V(x_{n+1}) \leq V(x_n)$, hence $\langle v_n \rangle_{n=0}^\infty$ is a decreasing sequence of real numbers and thus Cauchy. The inequality $d(x_n, x_m) \leq V(x_n) - V(x_m)$ for $m > n$ implies that $\langle x_n \rangle_{n=0}^\infty$ is also Cauchy, hence it has a limit, say x_∞ . By the continuity of f and the definition of x_n we see that $x_\infty = \lim_{n \rightarrow \infty} x_n = \lim_{n \rightarrow \infty} f(x_n) = f(\lim_{n \rightarrow \infty} x_n) = f(x_\infty)$, so indeed, x_∞ is a fixed point of f . \square

In the rest of this work we will show that all of these items reverse, except for (iii). (Note that the function in (iii) is required, by our definition of a Caristi system, to be Baire or Borel.)

Proposition 5.6. *The CFP for $\mathcal{X} = 2^\mathbb{N}$ and both f and V continuous implies WKL₀ over RCA₀.*

Proof. We prove the contrapositive over RCA₀. Assume that WKL₀ fails, and let $T \subseteq 2^{<\mathbb{N}}$ be an infinite binary tree with no infinite path. We define continuous $f: 2^\mathbb{N} \rightarrow 2^\mathbb{N}$ and $V: 2^\mathbb{N} \rightarrow [0, 3]$ so that $(2^\mathbb{N}, f, V)$ is a Caristi system with no fixed point. Our V is the potential from the proof that Ekeland's variational principle for continuous potentials on $2^\mathbb{N}$ implies WKL₀ of [4, Proposition 9.1].

As in the proof of [4, Proposition 9.1], let $T^\circ = \{\sigma \in T : \sigma \frown 0, \sigma \frown 1 \notin T\}$ be the set of leaves of T . The set T° is infinite because T is infinite but has no path. For each $\sigma \in T^\circ$, let

$$A_\sigma = \{i < |\sigma| : \neg(\exists \tau \in T)(|\tau| = |\sigma| + 1 \wedge \tau \sqsupseteq (\sigma \frown i) \wedge (1 - \sigma(i)))\}.$$

For each $\sigma \in 2^{<\mathbb{N}}$, let $\tilde{\sigma} \in 2^{<\mathbb{N}}$ be the sequence of length $2|\sigma|$ where $\tilde{\sigma}(2i) = 0$ and $\tilde{\sigma}(2i+1) = \sigma(i)$ for all $i < |\sigma|$. Define $\tilde{T} = \{\tilde{\sigma} : \sigma \in T\}$, $\tilde{T}^\circ = \{\tilde{\sigma} : \sigma \in T^\circ\}$, and

$$S = \{\tau \in 2^{<\mathbb{N}} : (\forall \sigma \in T)(|\sigma| \leq |\tau| \rightarrow \tau \not\sqsupseteq \tilde{\sigma}) \wedge (\exists \sigma \in T)(|\sigma| \leq |\tau| \wedge \tau \upharpoonright (|\tau| - 1) \sqsubset \tilde{\sigma})\}.$$

The set S consists of the shortest binary sequences that move away from \tilde{T} before reaching an element of \tilde{T}° . The elements of $\tilde{T}^\circ \cup S$ are pairwise incomparable, and the fact that T has no infinite path

implies that for every $x \in 2^{\mathbb{N}}$, there is a $\sigma \in \tilde{T}^\circ \cup S$ with $\sigma \sqsubseteq x$. Define the continuous function $V: 2^{\mathbb{N}} \rightarrow [0, 3]$ by

$$V(x) = \begin{cases} 2 - \sum_{i \in A_\sigma} 2^{-2i} & \text{if } x \sqsupseteq \tilde{\sigma} \text{ for a } \sigma \in T^\circ \\ 3 & \text{if } x \sqsupseteq \tau \text{ for a } \tau \in S. \end{cases}$$

One may obtain a code for V via [4, Lemmas 3.8 and 3.9].

Before defining f , observe that for any $\sigma \in T^\circ$, there is a $\tau \in T$ with $|\tau| = |\sigma| + 1$ because T is infinite. Such a τ does not extend σ because σ is a leaf, so there is an $i < |\sigma|$ such that $\tau \sqsupseteq (\sigma \upharpoonright i) \frown (1 - \sigma(i))$. Thus there is an $i < |\sigma|$ with $i \notin A_\sigma$. Given $\sigma \in T^\circ$, let i_σ be greatest such that $i_\sigma < |\sigma|$ and $i_\sigma \notin A_\sigma$, and let σ_+ be the first element of T° in lexicographic order with $|\sigma_+| > |\sigma|$ and $\sigma_+ \sqsupseteq (\sigma \upharpoonright i_\sigma) \frown (1 - \sigma(i_\sigma))$. Then $i_\sigma \in A_{\sigma_+}$ because any $\tau \in T$ with $\tau \sqsupseteq (\sigma_+ \upharpoonright i_\sigma) \frown (1 - \sigma_+(i_\sigma))$ satisfies $\tau \sqsupseteq \sigma \upharpoonright (i_\sigma + 1)$ and therefore also satisfies $|\tau| \leq |\sigma| < |\sigma_+|$ by the maximality of i_σ . Furthermore, if $i < i_\sigma$ and $i \in A_\sigma$, then $i \in A_{\sigma_+}$ as well because $\sigma_+ \upharpoonright i_\sigma = \sigma \upharpoonright i_\sigma$ and $|\sigma| < |\sigma_+|$.

Fix the first $\eta \in T^\circ$. Define the continuous function $f: 2^{\mathbb{N}} \rightarrow 2^{\mathbb{N}}$ following V by

$$f(x) = \begin{cases} \tilde{\sigma}_+ \frown 0^{\mathbb{N}} & \text{if } x \sqsupseteq \tilde{\sigma} \text{ for a } \sigma \in T^\circ \\ \tilde{\eta} \frown 0^{\mathbb{N}} & \text{if } x \sqsupseteq \tau \text{ for a } \tau \in S. \end{cases}$$

Again, one obtains a code for f via [4, Lemmas 3.8 and 3.9]. The function f has no fixed point. However, $(2^{\mathbb{N}}, f, V)$ is a Caristi system. Let $x \in 2^{\mathbb{N}}$. If $x \sqsupseteq \tilde{\sigma}$ for a $\sigma \in T^\circ$, then $d(x, f(x)) = 2^{-2i_\sigma - 1}$ and

$$\begin{aligned} V(x) - V(f(x)) &= \sum_{i \in A_{\sigma_+}} 2^{-2i} - \sum_{i \in A_\sigma} 2^{-2i} \geq 2^{-2i_\sigma} - \sum_{\substack{i \in A_\sigma \\ i > i_\sigma}} 2^{-2i} \\ &\geq 2^{-2i_\sigma} - 2^{-2i_\sigma - 1} = 2^{-2i_\sigma - 1} = d(x, f(x)), \end{aligned}$$

where the first inequality is because $i_\sigma \in A_{\sigma_+} \setminus A_\sigma$ and because $i \in A_\sigma \rightarrow i \in A_{\sigma_+}$ for $i < i_\sigma$. If instead $x \sqsupseteq \tau$ for a $\tau \in S$, then

$$V(x) - V(f(x)) = 1 + \sum_{i \in A_\eta} 2^{-2i} \geq d(x, f(x)).$$

Therefore $(2^{\mathbb{N}}, f, V)$ is a Caristi system with no fixed point, which completes the proof. \square

Peng and Yamazaki [9] showed that ACA_0 is equivalent to the CFP for continuous f and V . We sharpen this result by showing that we may assume that \mathcal{X} is the Baire space.

Proposition 5.7. *The CFP for $\mathcal{X} = \mathbb{N}^{\mathbb{N}}$ and both f and V continuous implies ACA_0 over RCA_0 .*

Proof. We prove the contrapositive over RCA_0 . Assume that ACA_0 fails, and let $h: \mathbb{N} \rightarrow \mathbb{N}$ be an injection whose range does not exist as a set. Let $T \subseteq \mathbb{N}^{<\mathbb{N}}$ be the tree constructed from h as in the proof that König's lemma implies ACA_0 of [13, Theorem III.7.2]. This T is the set of all $\sigma \in \mathbb{N}^{<\mathbb{N}}$ such that $(\forall m < |\sigma|)(\forall n < |\sigma|)(h(m) = n \leftrightarrow \sigma(n) = m + 1)$ and $(\forall n < |\sigma|)(\sigma(n) > 0 \rightarrow h(\sigma(n) - 1) = n)$. The tree T does not have an infinite path because the range of h does not exist as a set. We define continuous $f: \mathbb{N}^{\mathbb{N}} \rightarrow \mathbb{N}^{\mathbb{N}}$ and $V: \mathbb{N}^{\mathbb{N}} \rightarrow [0, 3]$ so that $(\mathbb{N}^{\mathbb{N}}, f, V)$ is a Caristi system with no fixed point.

To every $\sigma \in T$, assign a $\sigma_+ \in T$ with $|\sigma_+| > |\sigma|$ and $\{n < |\sigma_+| : \sigma_+(n) > 0\} \supseteq \{n < |\sigma| : \sigma(n) > 0\}$ as follows. Given $\sigma \in T$, let $k = \max\{|\sigma|, \max\{\sigma(n) : n < |\sigma|\}\}$. Let X be the finite set $X = \{h(m) : m \leq k\}$, and let σ_+ be the sequence of length $k + 1$ where for $n \leq k$

$$\sigma_+(n) = \begin{cases} m + 1 & \text{if } n \in X \text{ and } h(m) = n \\ 0 & \text{if } n \notin X. \end{cases}$$

Then $\sigma_+ \in T$. Furthermore, if $n < |\sigma|$ and $\sigma(n) > 0$, then $\sigma(n) \leq k$, so $h(\sigma(n) - 1) = n \in X$, so $\sigma_+(n) = \sigma(n) > 0$.

Define continuous $f: \mathbb{N}^{\mathbb{N}} \rightarrow \mathbb{N}^{\mathbb{N}}$ and $V: \mathbb{N}^{\mathbb{N}} \rightarrow [0, 3]$ as follows, using [4, Lemmas 3.8 and 3.9] to obtain the codes. Given $x \in \mathbb{N}^{\mathbb{N}}$, let $\sigma \sqsubseteq x$ be the longest initial segment with $\sigma \in T$, which exists because x is not a path through T . Let

$$V(x) = 1 + 2^{-|\sigma|+1} - \sum_{\substack{n < |\sigma| \\ \sigma(n) > 0}} 2^{-n} \qquad f(x) = \sigma_+ \frown 0^{\mathbb{N}}.$$

Notice that $f(x) \neq x$ because $\sigma_+ \in T$ but $x \upharpoonright |\sigma_+| \notin T$ as $|\sigma_+| > |\sigma|$. So f has no fixed point. To see that $d(x, f(x)) \leq V(x) - V(f(x))$, first observe that the longest initial segment $\tau \sqsubseteq f(x)$ with $\tau \in T$ has the form $\tau = \sigma_+ \frown 0^\ell$ for some ℓ because $\sigma_+ \in T$. Therefore

$$V(f(x)) = 1 + 2^{-|\tau|+1} - \sum_{\substack{n < |\tau| \\ \tau(n) > 0}} 2^{-n} \leq 1 + 2^{-|\sigma_+|+1} - \sum_{\substack{n < |\sigma_+| \\ \sigma_+(n) > 0}} 2^{-n}$$

because $|\sigma_+| \leq |\tau|$ and $\{n < |\sigma_+| : \sigma_+(n) > 0\} = \{n < |\tau| : \tau(n) > 0\}$. Thus

$$\begin{aligned} V(x) - V(f(x)) &\geq 2^{-|\sigma|+1} - 2^{-|\sigma_+|+1} + \sum_{\substack{n < |\sigma_+| \\ \sigma_+(n) > 0}} 2^{-n} - \sum_{\substack{n < |\sigma| \\ \sigma(n) > 0}} 2^{-n} \\ &\geq 2^{-|\sigma|} + \sum_{\substack{n < |\sigma_+| \\ \sigma_+(n) > 0}} 2^{-n} - \sum_{\substack{n < |\sigma| \\ \sigma(n) > 0}} 2^{-n}, \end{aligned}$$

where the second inequality is because $|\sigma| < |\sigma_+|$. If $\sigma \sqsubseteq \sigma_+$, then $d(x, f(x)) = 2^{-|\sigma|}$, so

$$d(x, f(x)) = 2^{-|\sigma|} \leq 2^{-|\sigma|} + \sum_{\substack{n < |\sigma_+| \\ \sigma_+(n) > 0}} 2^{-n} - \sum_{\substack{n < |\sigma| \\ \sigma(n) > 0}} 2^{-n} \leq V(x) - V(f(x)),$$

where the first inequality holds because $\{n < |\sigma_+| : \sigma_+(n) > 0\} \supseteq \{n < |\sigma| : \sigma(n) > 0\}$. If $\sigma \not\sqsubseteq \sigma_+$, then let $j < |\sigma|$ be least with $\sigma(j) \neq \sigma_+(j)$. It must be that $\sigma(j) = 0$ and $\sigma_+(j) > 0$ because if $\sigma(j) > 0$, then $\sigma_+(j) > 0$ as well, in which case $\sigma(j) = \sigma_+(j)$ as both σ and σ_+ are in T . Therefore

$$2^{-j} \leq \sum_{\substack{n < |\sigma_+| \\ \sigma_+(n) > 0}} 2^{-n} - \sum_{\substack{n < |\sigma| \\ \sigma(n) > 0}} 2^{-n}$$

because $\sigma_+(j) > 0$, $\sigma(j) = 0$, and $\{n < |\sigma_+| : \sigma_+(n) > 0\} \supseteq \{n < |\sigma| : \sigma(n) > 0\}$. Thus

$$d(x, f(x)) = 2^{-j} \leq 2^{-|\sigma|} + \sum_{\substack{n < |\sigma_+| \\ \sigma_+(n) > 0}} 2^{-n} - \sum_{\substack{n < |\sigma| \\ \sigma(n) > 0}} 2^{-n} \leq V(x) - V(f(x)).$$

So $d(x, f(x)) \leq V(x) - V(f(x))$ in both cases. Therefore $(\mathbb{N}^{\mathbb{N}}, f, V)$ is a Caristi system with no fixed point, which completes the proof. \square

Proposition 5.8. *The CFP for $[0, 1]$ with Baire class 1 f implies ACA_0 over RCA_0 .*

Proof. We work in RCA_0 and prove the contrapositive. Over RCA_0 , ACA_0 is equivalent to the monotone convergence theorem (see [13, Theorem III.2.2]). In fact, by inspecting the proof of [13, Theorem III.2.2], ACA_0 is equivalent to the statement “every strictly increasing sequence of rationals in $[0, 1]$ has a supremum.” Thus we let $\langle c_n \rangle_{n \in \mathbb{N}}$ be a strictly increasing sequence of rationals in $[0, 1]$ with no supremum, and we define a Caristi system $\langle [0, 1], f, V \rangle$ with f Baire class 1, but with no fixed points.

First we define V . Let $V_n: [0, 1] \rightarrow [0, 2]$ be a piecewise linear function such that $V_n(x) = 2$ for $x \leq c_n$, $V_n(x) = x$ for $x \geq c_{n+1}$, and V_n descends linearly on $[c_n, c_{n+1}]$. Let $V = \lim_{n \rightarrow \infty} V_n$. Clearly the functions V_n are continuous and increasing on n , so V is a potential and it is easy to see that

$$V(x) = \begin{cases} 2 & \text{if } \exists n(x < c_n) \\ x & \text{otherwise.} \end{cases}$$

Now we define f , show that $([0, 1], f, V)$ is a Caristi system, and show that f has no fixed points. Define a sequence $\langle f_n \rangle_{n \in \mathbb{N}}$ of piecewise-linear functions $f_n: [0, 1] \rightarrow [0, 1]$ as follows. For each n , first find the sequence $\langle q_i^n \rangle_{i \leq n+2}$, where

- $q_0^n = 1$;
- for each $i \leq n$, q_{i+1}^n is the rational in (c_n, q_i^n) with the least code;
- $q_{n+2}^n = c_n$.

Now define f_n so that

- for $i \leq n$, f_n is linear on $[q_{i+1}^n, q_i^n]$ with $f(q_{i+1}^n) = q_{i+2}^n$ and $f(q_i^n) = q_{i+1}^n$;
- f_n is linear on $[q_{n+2}^n, q_{n+1}^n]$ with $f(q_{n+2}^n) = 1$ and $f(q_{n+1}^n) = q_{n+2}^n$;
- f_n is constantly 1 on $[0, q_{n+2}^n]$.

We show, for every $x \in [0, 1]$, that $f(x) = \lim_{n \rightarrow \infty} f_n(x)$ exists, that $d(x, f(x)) \leq V(x) - V(f(x))$, and that $f(x) \neq x$. Let $x \in [0, 1]$. First suppose that there is an n_0 such that $x < c_{n_0}$. In this case, $f_n(x) = 1$ for all $n \geq n_0$, so $\lim_{n \rightarrow \infty} f_n(x) = 1$. Thus $f(x) = 1 \neq x$. Moreover, $d(x, f(x)) \leq 1 = 2 - 1 = V(x) - V(f(x))$.

Now suppose that $\forall n (c_n \leq x)$. As x is not the supremum of $\langle c_n \rangle_{n \in \mathbb{N}}$, there is a rational v such that $v \leq x$ and $\forall n (c_n \leq v)$. Thus by $\text{I}\Sigma_1^0$ in the guise of the Π_1^0 least number principle, there is such a v whose code is least. Similarly, there is a rational u whose code is least such that $u < v$ and $\forall n (c_n \leq u)$. Let Q be the finite set of rationals $\geq u$ whose codes are at most the code of u . Let $\langle q_i \rangle_{i \leq m}$ be the longest sequence of elements from $Q \cup \{1\}$ where $q_0 = 1$ and, for each $i < m$, q_{i+1} is the element of $[0, q_i] \cap Q$ with the least code. Observe that $q_m = u$ and $q_{m-1} = v$. Now, by the choice of u , let $n_0 \geq m$ be large enough so that $p \leq c_{n_0}$ for every rational $p \in [0, u)$ whose code is less than the code of u . Then $n \geq n_0$ implies that $(\forall i \leq m)(q_i^n = q_i)$. Thus, as $x \in [q_{m-1}, q_{m-2}]$, $n \geq n_0$ implies that $f_n(x) = \frac{q_{m-1} - q_m}{q_{m-2} - q_{m-1}}(x - q_{m-1}) + q_m$. So $f(x) = \lim_{n \rightarrow \infty} f_n(x) = \frac{q_{m-1} - q_m}{q_{m-2} - q_{m-1}}(x - q_{m-1}) + q_m$. We thus have that $q_m \leq f(x) < x$. Hence $\forall n (c_n \leq q_m \leq f(x))$ because $q_m = u$. Therefore $V(x) = x$ and $V(f(x)) = f(x)$, so $d(x, f(x)) = x - f(x) = V(x) - V(f(x))$.

We have now defined a potential $V: [0, 1] \rightarrow [0, \infty)$ and a Baire class 1 function $f: [0, 1] \rightarrow [0, 1]$ such that $([0, 1], f, V)$ is a Caristi system but such that f has no fixed points. This completes the proof. \square

Let us put together some of our results so far:

Theorem 5.9 (RCA₀). *The following are equivalent:*

- (a) WKL₀;
- (b) the CFP for continuous V and compact \mathcal{X} ;
- (c) the CFP for continuous f, V on the Cantor space.

Proof. That (a) implies (b) is Proposition 5.5, (c) is a special case of (b), and (c) implies (a) is Proposition 5.6. \square

Theorem 5.10 (RCA₀). *The following are equivalent:*

- (a) ACA₀;
- (b) the CFP for either f or V continuous;
- (c) the CFP for compact \mathcal{X} ;
- (d) the CFP for Baire class 1 f on $[0, 1]$;
- (e) the CFP for continuous f, V on the Baire space.

Proof. Proposition 5.5 shows that (a) implies both (b) and (c). Item (d) is a case of (c) and (e) is a case of (b). That (d) implies (a) is Proposition 5.8 and (e) implies (a) is Proposition 5.7. \square

We remark that the strength of the CFP for compact \mathcal{X} , continuous f , and arbitrary V is left open, although by the above, it must fall between WKL₀ and ACA₀.

6. CARISTI'S THEOREM AND THE TLPP

In this section we will show that the unrestricted CFP is equivalent to TLPP₀; recall that we defined the latter in Section 2.

We first prove Caristi's theorem in TLPP₀. We derive this from a 'relativized' version of Ekeland's variational principle provable in TLPP₀. The idea is that for a potential V , we can find a potential V' which is the same as V on hyperarithmetically definable points (with suitable parameters) and so that V' has a critical point. Below, we let \mathcal{X}_{\leq}^Y be the set of all points of \mathcal{X} that are Σ_{\leq}^Y .

Definition 6.1. Let \mathcal{X} be a complete separable metric space and V be a potential on \mathcal{X} . Let $Z \subseteq \mathbb{N}$. A Σ_{\leq}^Z relativization of V is a potential $V' = \lim_{n \rightarrow \infty} V'_n$ whose Baire code is arithmetical on some set W and such that for every $x \in \mathcal{X}_{\leq}^{W \oplus Z}$ and $n > 0$,

$$V'_n(x) = \inf_{y \in \mathcal{X}_{\leq}^{W \oplus Z}} (V(y) + nd(x, y)). \quad (1)$$

We call W the *parameter* of V' .

Roughly speaking, V' represents a "smoothed" version of V in which discontinuities at points of high complexity are forgotten.

Lemma 6.2 (TLPP₀). *For every potential V and every set Z , there is a Σ_{\prec}^Z relativization of V .*

Proof. Fix Z and V . Note that for any set W , the function $V'_{n-1}(x)$ defined by (1) is uniformly continuous (when defined), because for $x, x', y \in \mathcal{X}$, we have that

$$|(V(y) + nd(x, y)) - (V(y) + nd(x', y))| \leq nd(x, x'),$$

hence $|V'_n(x) - V'(x')| \leq nd(x, x')$. It is moreover evident that $V'_n(x) \leq V'_m(x)$ whenever $n \leq m$, given that $nd(x, y) \leq md(x, y)$ for all x, y . It remains to show that $\lim_{n \rightarrow \infty} V'_n$ has a Baire code as a sequence of continuous functions.

Let \mathcal{X} be a complete separable metric space, let $V: \mathcal{X} \rightarrow [0, \infty)$ be a potential, and let Z be any set. The property $c < V(y) + nd(a, y)$ is Σ_1^0 , hence using TLPP₀ in the form of Lemma 2.4 item 3, we obtain sets Δ and Y so that for $x \in X$, $(a, c, n) \in \Delta$ if and only if for all $y \in \mathcal{X}_{\prec}^{\Delta \oplus Y \oplus Z}$, $c < V(y) + nd(a, y)$. We set $W := \Delta \oplus Y$ and write \mathcal{X}' for $\mathcal{X}_{\prec}^{W \oplus Z}$.

From Δ , we may arithmetically define a code Υ_n for V'_n . Let us represent open balls in \mathbb{R} as intervals (c, d) . Then, enumerate $B_r(a)$ into $\Upsilon_n((c, d))$ if there are c', d' such that $c < c' < d' < d$ and for all $b \in B_r(a) \cap X$, $V'_n(b) \in (c', d')$. We check that Υ_n thus defined satisfies Definition 3.5 and codes the desired function V'_n .

For Item (i), it is immediate that if $(c_0, d_0) \subseteq (c_1, d_1)$ and $B_r(a)$ is in $\Upsilon_n((c_0, d_0))$ via (c', d') , then the same sub-interval witnesses that $B_r(a)$ is in $\Upsilon_n((c_1, d_1))$. For Item (ii), if $\Upsilon_n((c_0, d_0)) \cap \Upsilon_n((c_1, d_1)) \neq \emptyset$, since these sets are open and X is dense, we may find $a \in \Upsilon_n((c_0, d_0)) \cap \Upsilon_n((c_1, d_1)) \cap X$, which implies that $V'_n(a) \in (c_0, d_0) \cap (c_1, d_1)$, so the two intervals must intersect.

It remains to check that Υ_n indeed codes the desired function V'_n of (1). Let $x \in \mathcal{X}$ and write $x = \lim_{i \rightarrow \infty} x_i$, with $x_i \in X$. By uniform continuity, the value $V'_n(x)$ as given by (1) is $\lim_{i \rightarrow \infty} V'_n(x_i)$, which exists in ACA₀ as a real number. Now, suppose that (c, d) is an interval containing $V'_n(x)$, and let c', d' be such that $c < c' < V'_n(x) < d' < d$. Since V'_n is continuous, for small enough δ , we see that $V'_n[B_\delta(x)] \subseteq (c', d')$. Let $a \in X$ be such that $d(a, x) < \delta/2$, so that $x \in B_{\delta/2}(a) \subseteq B_\delta(x)$. Then $V'_n[B_{\delta/2}(a)] \subseteq (c', d')$, so $V'_n[B_{\delta/2}(a)]$ is enumerated in $\Upsilon_n((c, d))$ and $x \in \Upsilon_n((c, d))$. Since c, d were arbitrary, we see that x is in the domain of the coded function by Υ_n and $V'_n(x)$ is indeed the value assigned to x . \square

When looking at points of not-too-high complexity, we want V' to look essentially the same as V . The next lemma makes this precise.

Lemma 6.3. *Let V be a potential on \mathcal{X} , and let $V' = \lim_{n \rightarrow \infty} V'_n$ be a Σ_{\prec}^Z relativization of V . Then, $V(x) = V'(x)$ for every $x \in \mathcal{X}'$.*

Proof. Let $x \in \mathcal{X}'$. From the definition of V'_n , we see that we may instantiate y as x and obtain $V'_n(x) \leq V(x) + nd(x, x) = V(x)$, so $V'(x) \leq V(x)$.

To show that $V(x) \leq V'(x)$, we show that $V(x) \leq V'(x) + \varepsilon$ for all rational $\varepsilon > 0$. It suffices to show that for all $\varepsilon > 0$ there is an n such that $V(x) \leq V'_n(x) + \varepsilon$.

Since V is lsc, let δ be so that $d(x, y) < \delta$ implies $V(y) \geq V(x) - \varepsilon$. Choose n so that $n\delta > V(x)$. Then, $V(x) \leq V(y) + nd(x, y) + \varepsilon$ for every $y \in \mathcal{X}'$: if $d(x, y) < \delta$ this is because $V(x) \leq V(y) + \varepsilon$ by our choice of δ , otherwise $V(x) \leq n\delta \leq nd(x, y)$ by our choice of n . We conclude that $V(x) \leq V'_n(y) + \varepsilon$, as needed. \square

As mentioned, V' is meant to be a version of V where discontinuities of high complexity are removed. The following makes this precise.

Lemma 6.4. *Let $V: \mathcal{X} \rightarrow [0, \infty)$ be any potential and V' a Σ_{\prec}^Z -relativization of V . If $x \in \mathcal{X}$ and $\varepsilon > 0$, then there is $y \in \mathcal{X}'$ such that $d(x, y) < \varepsilon$ and $V'(y) < V'(x) + \varepsilon$.*

Proof. Assume toward a contradiction that $x \in \mathcal{X}$ and $\varepsilon > 0$ are such that $V'(y) \geq V'(x) + \varepsilon$ whenever $y \in \mathcal{X}'$ is such that $d(x, y) < \varepsilon$.

Choose $n > V'(x)/\varepsilon + 1$ and let $y \in \mathcal{X}'$ be arbitrary. If $d(x, y) < \varepsilon$,

$$V'(y) + nd(x, y) \geq V'(y) \geq V'(x) + \varepsilon,$$

while if $d(x, y) \geq \varepsilon$, we see by our choice of n that

$$V'(y) + nd(x, y) \geq n\varepsilon > (V'(x)/\varepsilon + 1)\varepsilon = V'(x) + \varepsilon.$$

But $V(y) = V'(y)$ by Lemma 6.3, so we conclude that $V(y) + nd(x, y) \geq V'(x) + \varepsilon$ for all $y \in \mathcal{X}'$. We obtain $V'_n(x) > V'(x) + \varepsilon$ by (1), contradicting $V'_n(x) \leq V'(x)$. \square

With this, we can show that V' is indeed enveloped.

Lemma 6.5 (RCA₀). *Let V be a potential on \mathcal{X} , and let $V' = \lim_{n \rightarrow \infty} V'_n$ be a $\Sigma^Z_{<}$ relativization of V . Then, $V'_n(x) = V'_{(n)}(x)$ for all n .*

Proof. Fix $x \in \mathcal{X}$. First we show for $y \in \mathcal{X}$ arbitrary that $V'_n(x) \leq V'(y) + nd(x, y)$. Let $\varepsilon > 0$. Using Lemma 6.4, let $y' \in \mathcal{X}'$ be such that $d(y, y') < \varepsilon/2(n+1)$ and $V'(y') < V'(y) + \varepsilon/2$. Then,

$$\begin{aligned} V'_n(x) &\leq V(y') + nd(x, y') && \text{by definition of } V'_n, \\ &= V'(y') + nd(x, y') && \text{since } y' \in \mathcal{X}', \\ &< V'(y) + nd(x, y) + \varepsilon && \text{by our choice of } y'. \end{aligned}$$

Since ε was arbitrary, $V'_n(x) \leq V'(y) + nd(x, y)$.

Finally we check that if $\varepsilon > 0$, then there is $y \in \mathcal{X}$ such that $V'_n(x) + \varepsilon > V'(y) + nd(x, y)$. By definition, $V'_n(x) = \inf_{y \in \mathcal{X}'} (V(y) + nd(x, y))$, so we can choose $y \in \mathcal{X}'$ such that $V'_n(x) + \varepsilon > V(y) + nd(x, y)$. Since $y \in \mathcal{X}'$, $V'(y) = V(y)$, so that

$$V'_n(x) + \varepsilon > V'(y) + nd(x, y),$$

as required. \square

Since ACA₀ proves that all enveloped potentials have a critical point by Theorem 5.2 and Lemma 5.3, we obtain the following.

Corollary 6.6 (ACA₀). *Every $\Sigma^Z_{<}$ relativization with parameter W has a critical point that is arithmetical in $W \oplus Z$ (hence $\Sigma^{W \oplus Z}_{<}$ if $|<|$ is infinite).*

With this, we may prove Caristi's theorem for Baire or Borel functions.

Proposition 6.7 (TLPP₀). *Caristi's fixed point theorem holds for arbitrary lower semi-continuous potentials V and arbitrary Baire or Borel functions f .*

Proof. Since TLPP₀ extends ATR₀, we may appeal to Lemma A.5 to see that every Baire function is Borel, so we may assume that f is Borel.

Let \mathcal{X} be a complete separable metric space, let $\Phi \subseteq \mathbb{N} \times X \times \mathbb{Q}^{>0} \times \mathbb{Q}$ code a lower semi-continuous potential on \mathcal{X} , and let f be a class-ot(\prec) Borel function coded by Υ ; it is convenient to assume that ot(\prec) is infinite. By Lemma 6.2, V has a $\Sigma^{\Upsilon}_{<}$ relativization V' with parameter W , and by Corollary 6.6, V' has a critical point x_* that is arithmetical in $W \oplus \Upsilon$. Since TLPP₀ extends ATR₀, we may use Lemma 3.6 to see that $f(x_*)$ is $\Sigma^{W \oplus \Upsilon}_{<}$. We have $d(x_*, f(x_*)) \leq V(x_*) - V(f(x_*))$, and since $V(f(x_*)) = V'(f(x_*))$, $V(x_*) = V'(x_*)$, and x_* is a critical point of V' , this implies that $x_* = f(x_*)$, as required. \square

Our next goal is to prove that the Caristi fixed point theorem for Baire/Borel functions is equivalent to TLPP₀ over RCA₀. As a first step, we show that ATR₀ is already provable at a rather low stage in the Baire hierarchy.

Theorem 6.8. *Over RCA₀, Caristi's theorem for Baire class-1 f implies ATR₀.*

Proof. Since Caristi's theorem for continuous functions implies ACA₀, we can work over ACA₀. Recall that the formula H describes the transfinite recursion for $\theta_*(n, Y)$ which defines a universal computably enumerable set relative to Y . Here, we may safely use a parameter $Y \in \mathbb{N}^{\mathbb{N}}$ instead of a member of $2^{\mathbb{N}}$. By the normal form theorem, we write $\theta_*(n, Y) \equiv \exists m \theta_0(m, n, Y \upharpoonright m)$.

In what follows, we will emulate the transfinite recursion on $\mathbb{N}^{\mathbb{N}}$. Let \prec be a well-order on \mathbb{N} . We will construct a sequence of trees $\langle T_\alpha : \alpha \in \mathbb{N} \rangle$ by Δ^0_1 -transfinite recursion which is implied from ACA₀ (see [2, 5]). For a given $\alpha \in \mathbb{N}$ and $\langle T_\beta \subseteq \mathbb{N}^{<\mathbb{N}} : \beta \prec \alpha \rangle$, we let $T^\alpha = \bigoplus_{\beta \prec \alpha} T_\beta := \{ \sigma \in \mathbb{N}^{<\mathbb{N}} : \text{for any } \beta < |\sigma|, \sigma_\beta \in T_\beta \text{ if } \beta \prec \alpha \text{ and } \sigma_\beta \text{ is a sequence of 0's otherwise} \}$, where $\sigma_i \in \mathbb{N}^{<\mathbb{N}}$ is $\sigma_i(x) = \sigma(\langle x, i \rangle)$ if $\langle x, i \rangle < |\sigma|$. (Here, the domain \mathbb{N} may be identified with $\mathbb{N} \times \mathbb{N}$ by the pairing function. Without loss of generality, we may assume that $\langle x, i \rangle \geq i$ in the sense of \mathbb{N} .)

Next we construct T_α from T^α . Here, the range \mathbb{N} may be identified with $\{0, 1\} \times \mathbb{N}^{<\mathbb{N}}$, and for a given $n = \langle s, \tau \rangle \in \mathbb{N}$, we write $s = (n)^\circ$ and $\tau = (n)^\circ$. (The idea here is that the second coordinate encodes a path $g \in [T^\alpha]$ and the first coordinate encodes the Turing jump of g .) Then we define T_α as follows: $\sigma \in T_\alpha$ if

- for any $s < |\sigma|$, $s \leq |(\sigma(s))^\circ|$ and $(\sigma(s))^\circ \in T^\alpha$,

- for any $t < s < |\sigma|$, $(\sigma(t))^\circ \sqsubset (\sigma(s))^\circ$,
- for any $s < |\sigma|$, $(\sigma(s))_\circ = 0 \rightarrow (\forall m \leq |(\sigma(|\sigma| - 1))^\circ|) \neg \theta_0(m, s, (\sigma(|\sigma| - 1))^\circ \upharpoonright m)$, and
- for any $s < |\sigma|$, $(\sigma(s))_\circ = 1 \rightarrow (\exists m \leq |(\sigma(s))^\circ|) \theta_0(m, s, (\sigma(s))^\circ \upharpoonright m)$.

Now, let $\langle T_\alpha \subseteq \mathbb{N}^{<\mathbb{N}} : \alpha \in \mathbb{N} \rangle$ be the result of the transfinite recursion. If $g \in [T_\alpha]$ for some $\alpha \in \mathbb{N}$, we let $(g)^\circ := \bigcup \{(g(s))^\circ : s \in \mathbb{N}\}$ and $(g)_\circ := \{g : (g(s))_\circ = 1\}$, then $(g)^\circ \in T_\alpha$ and $(g)_\circ = \{n : \theta_*(n, (g)^\circ)\}$. Moreover, let $f_\beta := ((g)^\circ)_\beta = \bigcup \{(g(s))^\circ_\beta : s \in \mathbb{N}\}$ for $\beta \prec \alpha$, and $f_\alpha = g$. then, by the definition, we have $f_\beta \in [T_\beta]$ for any $\beta \prec \alpha$. Now, by arithmetical transfinite induction up to α , we may verify that f_β computes a set Y^β such that $H_\theta(\prec_\beta, Y^\beta)$ for any $\beta \prec \alpha$, where \prec_γ is a restriction of \prec to the domain $\{\gamma' \in \mathbb{N} : \gamma' \prec \gamma\}$. Hence, if $\hat{f} \in [\bigoplus_{\alpha \in \mathbb{N}} T_\alpha]$, then $(\hat{f})_\alpha \in [T_\alpha]$ for any $\alpha \in \mathbb{N}$, and thus \hat{f} computes a set Y such that $H(\prec, Y)$.

Finally, we construct a Caristi system whose fixed point is a path of $\bigoplus_{\alpha \in \mathbb{N}} T_\alpha$. We define a potential function $V : \mathbb{N}^\mathbb{N} \rightarrow [0, \infty)$ by $V(f) = \sum \{2^{-\alpha} : (f)_\alpha \notin [T_\alpha], \alpha \in \mathbb{N}\}$; this defines a total potential on $\mathbb{N}^\mathbb{N}$ provably in ACA_0 ([4], together with Lemma A.4). We will construct a Baire class 1 function $F : \mathbb{N}^\mathbb{N} \rightarrow \mathbb{N}^\mathbb{N}$ which ‘descends along’ V . For each $s \in \mathbb{N}$, we will define a continuous function $F_s : \mathbb{N}^\mathbb{N} \rightarrow \mathbb{N}^\mathbb{N}$ as follows. For a given $f \in \mathbb{N}^\mathbb{N}$, let $I_{f,s} := \{\alpha < s : (f \upharpoonright s)_\alpha \notin T_\alpha, \alpha \in \mathbb{N}\}$. If $I_{f,s} = \emptyset$, then put $F_s(f) = f$. Otherwise, let β_s be the \prec -smallest element of $I_{f,s}$. Put $(F_s(f))_\gamma = (f)_\gamma$ if $\gamma \neq \beta_s$, and $(F_s(f))_{\beta_s}$ to be the s -approximation of a path of T_{β_s} computed from (the s -approximation of) $f^{\beta_s} := \bigoplus_{\gamma \prec \beta_s} f_\gamma$. (Note that $f^{\beta_s} \upharpoonright s \in T^{\beta_s}$ by the definition.) More formally, we define $h = (F_s(f))_{\beta_s}$ as follows: for $t < s$ define $h(t)$ inductively as $h(t) = (0, f^{\beta_s} \upharpoonright \bar{t})$ if $(\forall m \leq s) \neg \theta_0(m, t, f^{\beta_s} \upharpoonright m)$ and $\bar{t} = \max\{t\} \cup \{|(h(t'))^\circ| : t' < t\}$ and $h(t) = (1, f^{\beta_s} \upharpoonright \bar{t})$ if \bar{t} is the smallest $u \leq s$ such that $(\exists m \leq u) \theta_0(m, t, f^{\beta_s} \upharpoonright m)$ and $u \geq \max\{t\} \cup \{|(h(t'))^\circ| : t' < t\}$, and put $h(t) = 0$ for $t \geq s$. It is a routine to check that such F_s is continuous.

If $f \in [\bigoplus_{\alpha \in \mathbb{N}} T_\alpha]$, then $F_s(f) = f$ for any $s \in \mathbb{N}$. Otherwise, let $\hat{\beta}$ be the \prec -smallest α such that $(f)_\alpha \notin [T_\alpha]$. Then, for large enough s , $\beta_s = \hat{\beta}$. Thus, $g = \lim_{s \rightarrow \infty} F_s(f)$ exists and $(g)_\gamma = (f)_\gamma$ for any $\gamma \neq \hat{\beta}$ and $(g)_{\hat{\beta}} \in [T_{\hat{\beta}}]$. Moreover, we have $d(f, g) \leq 2^{-\hat{\beta}} \leq V(f) - V(g)$. Therefore, $F = \lim_{s \rightarrow \infty} F_s$ exists, $(\mathbb{N}^\mathbb{N}, F, V)$ forms a Caristi system and its fixed point is a path of $\bigoplus_{\alpha \in \mathbb{N}} T_\alpha$. \square

Question 6.9. Is Caristi’s theorem for Baire class-1 f provable in ATR_0 ?

Theorem 6.10 (RCA_0). *The following are equivalent:*

- (1) TLPP_0 ,
- (2) CFP for Baire f ,
- (3) CFP for Borel f , and
- (4) CFP for Borel f on the Baire space.

Proof. By Proposition 6.7, the first item implies the third. The first moreover implies the second, since TLPP_0 implies ATR_0 and so by Lemma A.5 shows that every Baire function is a Borel function. To see that the second implies the fourth, Theorem 6.8 shows that it implies ATR_0 , and therefore by Lemma A.7, it implies that every Borel function from $\mathbb{N}^\mathbb{N}$ to itself is a Baire function, so the fourth follows.

To complete the loop, it remains to show that the fourth implies the first. By Theorems 5.10 and 6.8, we may argue within ATR_0 . Using Lemma 2.4, we will show the equivalent form: given a sequence of trees $\langle T_n \rangle_{n \in \mathbb{N}}$ and a well-order \prec , we will obtain a pair of sets Z_0, Z_1 so that $n \in Z_0$ iff there is a path through T_n which is $\Sigma_{\prec}^{Z_0 \oplus Z_1 \oplus \langle T_n \rangle_{n \in \mathbb{N}}}$. To show this, we define $V : \mathbb{N}^\mathbb{N} \rightarrow [0, \infty)$ by $V(f) = \sum \{2^{-i} : (f)_i \notin [T_i]\}$ as in the proof of Theorem 6.8.

We wish to define a Borel function $F : \mathbb{N}^\mathbb{N} \rightarrow \mathbb{N}^\mathbb{N}$ by having $F(f)$ look for a g which is a counterexample to f being a critical point of V and which is not too much more complicated than f . Let \prec' be the successor of \prec . Then pick a uniform enumeration $\psi_0, \psi_1, \psi_2, \dots$ of the partial computable functions and, given f , we let Y_f be the set such that $H(\prec', Y_f, f \oplus \langle T_n \rangle_{n \in \mathbb{N}})$. Then for each e such that $\psi_e^{Y_f}$ is total, we can let $g_e = \psi_e^{Y_f}$. We wish to choose $F(f)$ to be g_e where e is least such that $0 < d(f, g_e) \leq V(f) - V(g_e)$ if there is such an e , and to be f otherwise.

We need to check that F is Borel. It suffices to show that, for any finite sequence $\sigma \in \mathbb{N}^{<\mathbb{N}}$, the set of f such that there is e such that $\psi_e^{Y_f}$ exists and satisfies $\sigma \sqsubset g_e$ and $0 < d(f, g_e) \leq V(f) - V(g_e)$ is Borel.

The main step is translating the construction of Y_f into a statement about Borel sets. By recursion on $\alpha \in |\prec'|$ we argue that the set of f such that $m \in (Y_f)_\alpha$ is Borel: when α is minimal, $m \in (Y_f)_\alpha$ iff

$\theta_*(m, \emptyset, f \oplus \langle T_n \rangle_{n \in \mathbb{N}})$, and since θ_* is Σ_1^0 , the set of such f is open, namely a union of those initial segments that witness this.

Suppose that, for all m , the set of f such that $m \in (Y_f)_{<\alpha}$ is Borel. Then $m \in (Y_f)_\alpha$ if and only if $\theta_*(m, (Y_f)_{<\alpha}, f \oplus \langle T_n \rangle_{n \in \mathbb{N}})$. This is a union of sets which are, by the recursion, Borel sets intersected with open sets, and is therefore also a Borel set. In particular, for each e , the set of f such that $\psi_e^{Y_f}$ is total is a Borel set.

The set of f such that $V(f) < q$ is easily seen to be Borel. Since, for any i and j , the set of f such that $\psi_e^{Y_f}(i) = j$ is Borel, also for any rational q , the set of f such that $0 < d(f, g_e) < q$ is Borel, as is the set of f such that $V(g_e) < q$. Then the set of f such that $0 < d(f, g_e) \leq V(f) - V(g_e)$ is precisely the set of f such that, for every pair q, q' such that $V(f) < q$ and $V(g_e) \geq q'$, $0 < d(f, g_e) < q - q'$, which is also Borel.

Therefore the function F is Borel. By CFP for Borel functions, there is a fixed point f_* for F , and so $F(f_*) = f_*$. Let $X = \{\beta : (f_*)_\beta \in [T_\beta]\}$. Clearly if $\beta \in X$ then T_β has a path. Conversely, suppose that $\beta \notin X$, and suppose for a contradiction that T_β has a path h which is $\Sigma_{<\omega}^{X \oplus f_* \oplus \langle T_n \rangle_{n \in \mathbb{N}}}$. Then we may define $g(\langle \gamma, n \rangle)$ by $h(n)$ if $\gamma = \beta$ and $f_*(\langle \gamma, n \rangle)$ otherwise, so g is $\Sigma_{<\omega}^{X \oplus f_* \oplus \langle T_n \rangle_{n \in \mathbb{N}}}$, and therefore $\Sigma_{<\omega}^{f_* \oplus \langle T_n \rangle_{n \in \mathbb{N}}}$. It follows that $g = g_e$ for some e .

Moreover, $0 < d(f_*, g) \leq 2^{-(\beta, 0)} \leq 2^{-\beta} = V(f_*) - V(g)$, which means that $F(f_*) = g_{e_0}$ for the least e_0 with $0 < d(f_*, g_{e_0}) \leq V(f_*) - V(g_{e_0})$, and so we must have $F(f_*) \neq f_*$, which is a contradiction. Therefore X and f_* are the necessary witnesses. \square

APPENDIX A. REMARKS ON FUNCTION CODES

In order to maintain some generality in the formalization of Caristi's theorem, we have based our presentation on Baire and Borel codes. In this Appendix we establish how these codes relate to each other, as well as to other codes used in the literature.

First, we recall the standard coding of continuous functions used in e.g. [13].

Definition A.1 (RCA_0 ; [13, Definition II.6.1]). Let $\mathcal{X} = \widehat{X}$ and $\mathcal{Y} = \widehat{Y}$ be complete separable metric spaces. A *continuous partial function* $f: \mathcal{X} \rightarrow \mathcal{Y}$ is coded by a set $\Phi \subseteq \mathbb{N} \times X \times \mathbb{Q}^{>0} \times Y \times \mathbb{Q}^{>0}$ that satisfies the properties below. Let us write $B_r(a) \xrightarrow{\Phi} B_q(b)$ for $\exists n(\langle n, a, r, b, q \rangle \in \Phi)$. Then, for all $a, a' \in X$, all $q, q' \in \mathbb{Q}$, and all $r, r' \in \mathbb{Q}^{>0}$, Φ should satisfy:

- (CF 1) if $B_r(a) \xrightarrow{\Phi} B_q(b)$ and $B_r(a) \xrightarrow{\Phi} B_{q'}(b')$, then $d(b, b') \leq q + q'$;
- (CF 2) if $B_r(a) \xrightarrow{\Phi} B_q(b)$ and $B_{r'}(a') \subseteq B_r(a)$, then $B_{r'}(a') \xrightarrow{\Phi} B_q(b)$;
- (CF 3) if $B_r(a) \xrightarrow{\Phi} B_q(b)$ and $B_q(b) \subseteq B_{q'}(b')$, then $B_r(a) \xrightarrow{\Phi} B_{q'}(b')$.

A point $x \in \mathcal{X}$ is in the domain of the function f coded by Φ if, for every $\varepsilon > 0$, there are $B_r(a) \xrightarrow{\Phi} B_s(b)$ such that $d(x, a) < r$ and $s < \varepsilon$. If $x \in \text{dom}(f)$, we define the value $f(x)$ to be the unique point $y \in \mathcal{Y}$ such that $d(y, b) \leq s$ for all $B_r(a) \xrightarrow{\Phi} B_s(b)$ with $d(x, a) < r$.

Lemma A.2 (RCA_0). *Let \mathcal{X}, \mathcal{Y} be metric spaces. A function $f: \mathcal{X} \rightarrow \mathcal{Y}$ can be coded as a continuous function in the sense of Definition A.1 if and only if it can be coded as a Borel function where the preimage of every open ball is open.*

Proof sketch. If Φ is a code for a continuous function f and $B_s(b)$ is any open ball in \mathcal{Y} , we can enumerate $f^{-1}[B_s(b)]$ by enumerating $B_r(a)$ if $B_r(a) \xrightarrow{\Phi} B_{s'}(b')$, where $d_{\mathcal{Y}}(b, b') + s' < s$ (the latter is needed since $B_r(a) \xrightarrow{\Phi} B_{s'}(b')$ only guarantees $B_r(a) \subseteq f^{-1}[\overline{B_s(b)}]$). Conversely, if Ψ is a Borel code for f , we note that for $B_s(b) \subseteq \mathcal{Y}$, $f^{-1}[B_s(b)]$ is represented in the form $\bigcup_{i \in \mathbb{N}} B_{r_i}(a_i)$. We thus enumerate $B_r(a) \xrightarrow{\Phi} B_s(b)$ if $a = a_i$ and $r = r_i$ for some i . \square

Indeed, the proof of Lemma A.2 is effective in the sense that there exist (provably in RCA_0) Δ_1^0 formulas which define Turing functionals for these conversions of codes, and thus any sequence of codes of functions in one way can be converted to the sequence of codes in the other way.

The results of [4] were originally stated with respect to the following coding of lower semi-continuous functions, but as we will see, it is equivalent to our Baire representation for them.

Definition A.3 (RCA_0 ; [4, Definition 4.1]). Let \mathcal{X} be a complete separable metric space. A *lower semi-continuous partial function* $V: \mathcal{X} \rightarrow \mathbb{R}$ is coded by a set $\Psi \subseteq \mathbb{N} \times X \times \mathbb{Q}^{>0} \times \mathbb{Q}$ that satisfies the

properties below. Let $B_r(a) \xrightarrow{\Psi} q$ denote $\exists n(\langle n, a, r, q \rangle \in \Psi)$. Then Ψ must satisfy that for all $a, a' \in X$, all $q, q' \in \mathbb{Q}$, and all $r, r' \in \mathbb{Q}^{>0}$,

(LSC 1) if $B_r(a) \xrightarrow{\Psi} q$ and $B_{r'}(a') \in B_r(a)$, then $B_{r'}(a') \xrightarrow{\Psi} q$, and

(LSC 2) if $B_r(a) \xrightarrow{\Psi} q$ and $q' < q$, then $B_r(a) \xrightarrow{\Psi} q'$.

A point $x \in \mathcal{X}$ is in the domain of the function V coded by Ψ if

$$y = \sup\{q \in \mathbb{Q} : (\exists \langle a, r \rangle \in X \times \mathbb{Q}^{>0})(B_r(a) \xrightarrow{\Psi} q \wedge d(x, a) < r)\}$$

exists, in which case $V(x) = y$. If V has codomain $[0, \infty)$ (in the sense that $B_r(a) \xrightarrow{\Psi} 0$ for every a, r), we call V a *potential*.

Let \mathcal{X} be a complete separable metric space. The idea behind Definition A.3 is that Ψ enumerates pairs $\langle B_r(a), [q, \infty) \rangle$ with the property that if V is the function being coded by Ψ and x is in $B_r(a) \cap \text{dom}(V)$, then $V(x)$ is in $[q, \infty)$.

Lemma A.4.

- (1) Over RCA_0 , it is provable that a function f has an lsc code as given by Definition A.3 if and only if it is a pointwise increasing limit of continuous functions.
- (2) Over $\Pi_1^1\text{-CA}_0$, it is provable that for every potential f and every $\alpha > 0$, the α -envelope of f exists.

Proof sketch. The second claim is proven in [4], so we focus on the first. First we approximate the indicator function of an open ball, $\chi_{B_r(a)}$; for $\varepsilon \in (0, 1)$, define $\chi_\varepsilon(x)$ to be 1 if $x \in B_{r(1-\varepsilon)}(a)$, 0 if $x \notin B_r(a)$, and otherwise $\chi_\varepsilon(x) = 1/\varepsilon - d(a, x)/\varepsilon r$. Then, if we define $\chi_{B_r(a)}^n := \chi_{2^{-n}r}$, it is clear that $\chi_{B_r(a)}^n \rightarrow \chi_{B_r(a)}$ as $n \rightarrow \infty$. If $\langle \langle B_i, q_i \rangle : i < m \rangle$ is a tuple of pairs consisting of an open ball and a positive rational, we may similarly approximate $\max_{i < m} q_i \chi_{B_i}$ by $\max_{i < m} q_i \chi_{B_i}^n$. If we enumerate an lsc code Ψ as $\{\langle B_i, q_i \rangle : i \in \mathbb{N}\}$, we may then approximate the function V coded by Ψ by diagonally approximating V as $\max_{i < n} q_i \chi_{B_i}^n$.

Conversely, if $V = \lim_{n \rightarrow \infty} V_n$ where $\langle V_n : n \in \mathbb{N} \rangle$ is pointwise increasing, we may define a code for V by putting $B \xrightarrow{\Psi} q$ if there is n such that $V_n(x) > q$ for all $x \in B$, which may be extracted from the continuous code for V_n . \square

Lemma A.5 (ATR_0). *Every Baire function is Borel.*

Proof. Let $f = f_\Xi$ be a Baire function coded by Ξ . It suffices to show that for all $B_r(a)$, the inverse image $f^{-1}[B_r(a)]$ is a Borel set in a uniform way. We show this by recursion on Ξ . The case where f is continuous is given by Lemma A.2, so we assume otherwise.

Suppose we have $f = \lim_{n \rightarrow \infty} f_n$ and each f_n is given as a Borel function. Then we may set

$$f^{-1}[B_r(a)] = \bigcup_{r' < r} \bigcup_n \bigcap_{m \geq n} f_m^{-1}[B_{r'}(a)].$$

\square

In general it is not true that every Borel function is Baire, but for many specific spaces this does hold. However, it does hold for *zero-dimensional spaces*, are spaces that have a basis consisting of clopen balls.

Lemma A.6 (ATR_0). *If \mathcal{X} is zero-dimensional, then every Borel function from \mathcal{X} to \mathbb{N} is Baire.*

Before beginning the proof, it will be helpful to refine our definitions a bit. Let \mathcal{A} be the algebra generated by the basic clopen sets of \mathcal{X} —that is, the collection of sets generated from basic clopen sets by complements, finite unions, and finite intersections. We can choose some encoding of the elements of \mathcal{A} by natural numbers.

We next modify our notion of a Borel code slightly. First, we restrict ourselves to codes where the levels alternate between unions and intersections—that is, which have the form $\bigcup_n \bigcap_m \bigcup_p \dots$. We can easily obtain a code with this property from one without by compressing runs of numbers with the same parity. For example, when $\sigma^\frown \langle 2n \rangle \in S$ is not a leaf, we can replace this node with nodes of the form

$$\sigma^\frown \langle 2 \langle 2n, 2m_1, \dots, 2m_k \rangle \rangle$$

such that $\sigma \frown \langle 2n, 2m_1, \dots, 2m_k \rangle \in S$, and similarly converting runs of odd numbers to a single odd number.

Second, we require that the top level be a union—that is, unless S contains only a single leaf, we require that the unique n so that $\langle n \rangle \in S$ be odd. This is easily arranged, because if S does not have this property then we replace it with $\{\langle 1 \rangle \frown \sigma : \sigma \in S\}$.

Finally, we allow leaves to be labeled by elements of \mathcal{A} ; this does not change what sets are Borel, though it can slightly reduce the complexity. For purposes of the proof, we call these *clean Borel codes*. We note that, because leaves are labeled by elements of \mathcal{A} , a clean Borel code is not quite a Borel code. Nonetheless, the basic properties of Borel codes hold for clean Borel codes with no changes to the proofs.

The advantage to clean Borel codes is that we can easily take finite unions and intersections without changing the complexity, and can take the complement while increasing the complexity by at most one (because we need to add an extra, trivial, union step at the root).

Proof of Lemma A.6. We assume f is given as a clean Borel function—that is, the code for f gives us, for each n , a clean Borel code for $f^{-1}(n)$.

We will define a well-founded tree U and, for each node $\sigma \in U$, a clean Borel function f_σ . Additionally, we need to give some sort of ordinal bound on the Borel codes we use, which will ensure that they are getting simpler as we progress to larger nodes in U ; we take an approach which is inefficient but relatively simple to describe. Take the tree combining all the Borel codes for f —that is, the tree Υ with branches indexed by \mathbb{N} , and above each n , the Borel code S^n for $f^{-1}(n)$. Then Υ is a well-founded tree.

For each σ and each n , we will have an assignment π_σ^n from the tree S_σ^n encoding $f_\sigma^{-1}(n)$ to Υ so that if $\tau \sqsubset \tau'$ then $\pi_\sigma^n(\tau') <_{\text{KB}(\Upsilon)} \pi_\sigma^n(\tau)$. We have $\langle \rangle \in U$ and $f_{\langle \rangle} = f$. The map $\pi_{\langle \rangle}^n$ is the inclusion of S^n in Υ .

If, for every n , $f_\sigma^{-1}(n)$ is presented as an element of \mathcal{A} (that is, the clean Borel code for $f_\sigma^{-1}(n)$ is simply a single leaf labeled by an element of \mathcal{A}) then σ is a leaf of U .

Otherwise, we will define a sequence of functions $f_{\sigma \frown \langle s \rangle}$ so that $\lim_{s \rightarrow \infty} f_{\sigma \frown \langle s \rangle} = f_\sigma$. If any $f_\sigma^{-1}(n) = B \in \mathcal{A}$, we may replace it with $f_\sigma^{-1}(n) = \bigcup_{i=1}^1 B$, so we assume that, for all n , $f_\sigma^{-1}(n) = \bigcup_i S_i^n$. Suppose that either $\{\pi_\sigma^n(\langle \rangle) : n \in \mathbb{N}\}$ doesn't have a largest element or $S_i^n \in \mathcal{A}$ for any $i, n \in \mathbb{N}$. Then we define

$$f_{\sigma \frown \langle s \rangle}^{-1}(n) = \bigcup_{i \leq s} S_i^n$$

for $n < s$,

$$f_{\sigma \frown \langle s \rangle}^{-1}(s) = \bigcap_{n < s} \bigcap_{i \leq s} \overline{S_i^n}$$

and $f_{\sigma \frown \langle s \rangle}^{-1}(n) = \emptyset$ for $n > s$. Then it is immediate that $\lim_{s \rightarrow \infty} f_{\sigma \frown \langle s \rangle} = f_\sigma$. The functions $\pi_{\sigma \frown \langle s \rangle}^n$ can be defined in the obvious way by copying over the definitions of the π_σ^n where possible and, in $\pi_{\sigma \frown \langle s \rangle}^n$, mapping the extra levels to successors. In particular, $\sup_n \pi_{\sigma \frown \langle s \rangle}^n(\langle \rangle) < \sup_n \pi_\sigma^n(\langle \rangle)$.

Otherwise, $\{\pi_\sigma^n(\langle \rangle) : n \in \mathbb{N}\}$ has a largest element and at least one $f_\sigma^{-1}(n)$ has the form $\bigcup_i \bigcap_j S_{i,j}^n$; then, again, we may assume that every $f_\sigma^{-1}(n)$ has this form (by replacing $\bigcup_i B_i$ with $\bigcup_i \bigcap_{j=1}^1 B_i$). For each s and n , we set

$$f_{\sigma \frown \langle s \rangle}^{-1}(n) = \bigcup_{t_1} \left[\bigcap_{m \leq s} S_{t_1, m}^n \cap \bigcap_{\langle u_0, u_1 \rangle < \langle n, t_1 \rangle} \bigcup_{m \leq s} \overline{S_{u_1, m}^{u_0}} \right].$$

That is, $f_{\sigma \frown \langle s \rangle}(x)$ is chosen by finding the smallest pair $\langle n, t_1 \rangle$ such that $x \in \bigcap_{m \leq s} S_{t_1, m}^n$. Such a pair $\langle n, t_1 \rangle$ clearly exists and is unique, so $f_{\sigma \frown \langle s \rangle}$ also defines a unique Borel function. Since, for each x , there is an n so that, for some t_1 , $x \in \bigcap_m S_{t_1, m}^n$, there is some large enough finite s so that $s \notin \bigcup_{\langle u_0, u_1 \rangle < \langle n, t_1 \rangle} \bigcap_{m \leq s} \overline{S_{u_1, m}^{u_0}}$, and therefore $f_{\sigma \frown \langle s \rangle}(x) = f_\sigma(x)$; in particular, the functions $f_{\sigma \frown \langle s \rangle}$ converge pointwise to f_σ .

We need to define the functions $\pi_{\sigma \frown \langle s \rangle}^n$, and we have to do this a bit carefully to make sure that $\sup_n \pi_{\sigma \frown \langle s \rangle}^n(\langle \rangle) < \sup_n \pi_\sigma^n(\langle \rangle)$. For simplicity, assume that each $S_{t_1, m}^n = \bigcup_j S_{t_1, m, j}^n$. (In the cases where

this fails, $S_{t_1, m}^n \in \mathcal{A}$, and the coding is similar but simpler.) Then we have, for each t_1 ,

$$\bigcap_{m \leq s} S_{t_1, m}^n \cap \bigcap_{\langle u_0, u_1 \rangle < \langle n, t_1 \rangle} \bigcup_{m \leq s} \overline{S_{u_1, m}^{u_0}} = \bigcup_{j_1, \dots, j_s} \left[\bigcap_{m \leq s} S_{t_1, m, j_m}^n \cap \bigcap_{\langle u_0, u_1 \rangle < \langle n, t_1 \rangle} \bigcup_{m \leq s} \overline{S_{u_1, m}^{u_0}} \right],$$

and using the closure of clean Borel codes under finite unions and intersections, the interior of the union is a single clean Borel code. Note that π_σ^n maps the node corresponding to each $S_{u_1, m}^{u_0}$ (or $S_{t_1, m}^n$) to a level at least three nodes below $\sup_n \pi_{\sigma \frown \langle s \rangle}^n(\langle \rangle)$ (there must be a node above for \bigcap_j , then one for \bigcup_i , then one for the root of the Borel code). By mapping to level-wise maxima, we may arrange for $\pi_{\sigma \frown \langle s \rangle}^n$ to map the root of this intersection to a level at least three below $\sup_n \pi_{\sigma \frown \langle s \rangle}^n(\langle \rangle)$ as well. Then we can map the union over t_1, j_1, \dots, j_s to the level immediately above that, and the root to the level above that. In particular, $\pi_{\sigma \frown \langle s \rangle}^n(\langle \rangle) + 1 \leq \sup_n \pi_\sigma^n(\langle \rangle)$, and therefore $\sup_n \pi_{\sigma \frown \langle s \rangle}^n(\langle \rangle) < \sup_n \pi_\sigma^n(\langle \rangle)$.

Since $\text{KB}(\Upsilon)$ is well-ordered, the map from U to Υ given by $\sigma \mapsto \sup_n \pi_\sigma^n(\langle \rangle)$ shows that U is well-founded. The leaves σ of U are functions where each $f_\sigma^{-1}(n) \in \mathcal{A}$, and since each open ball is clopen, each element of \mathcal{A} is open, so f_σ^{-1} is continuous. We may replace each leaf with a code for f_σ as a continuous function, and we have therefore obtained a representation of f as a Baire function. \square

Lemma A.7 (ATR₀). *Every Borel function from $\mathbb{N}^{\mathbb{N}}$ to itself is Baire.*

Proof. First, given a Borel function f coded by Υ , let $f' : \mathbb{N}^{\mathbb{N}} \times \mathbb{N} \rightarrow \mathbb{N}$ be the function which maps $\langle \Lambda, n \rangle$ to the n -th position of $f(\Lambda)$. We view $\mathbb{N}^{\mathbb{N}} \times \mathbb{N}$ as a metric space in which $d(\langle \Lambda, n \rangle, \langle \Lambda', n' \rangle) = 2$ iff $n \neq n'$ and $d(\langle \Lambda, n \rangle, \langle \Lambda', n \rangle) = d(\Lambda, \Lambda') \leq 1$. Then f' is also Borel: the inverse image of a is the union over all sets $U_{\Upsilon_{\langle \sigma, 2^{-|\sigma|} \rangle}} \times \{n\}$, where $n \in \mathbb{N}$ and $\sigma \in \mathbb{N}^{<\mathbb{N}}$ such that $\sigma(n) = a$.

By the previous lemma, f' is also Baire. Taking the (continuous) leaves $f'_\sigma : \mathbb{N}^{\mathbb{N}} \times \mathbb{N} \rightarrow \mathbb{N}$, we can define $f_\sigma : \mathbb{N}^{\mathbb{N}} \rightarrow \mathbb{N}^{\mathbb{N}}$ by $f_\sigma(\Lambda) = \langle f'_\sigma(\langle \Lambda, n \rangle) : n \in \mathbb{N} \rangle$. This function is still continuous and respects the limits of the f'_σ sequence, so the same tree gives a representation of f as a Baire function. \square

FUNDING

Fernández-Duque was supported by the FWO-FWF Lead Agency grant G030620N (FWO)/I4513N (FWF). Shafer was supported by the *Fonds voor Wetenschappelijk Onderzoek – Vlaanderen* Pegasus program, the John Templeton Foundation grant ID 60842 *A new dawn of intuitionism: mathematical and philosophical advances* and by EPSRC grant EP/T031476/1 *Reverse mathematics of general topology*. The opinions expressed in this work are those of the authors and do not necessarily reflect the views of the John Templeton Foundation. Towsner was partially supported by NSF grant DMS-2054379. Yokoyama was partially supported by JSPS KAKENHI grant number 19K03601 and 21KK0045.

ACKNOWLEDGMENTS

David Fernández-Duque would like to thank Carlos Bosch Giral, César Luis García García, Claudia Gómez Wulschner, Rigoberto Vera Mendoza, and the other attendees of the ITAM Analysis seminar, for introducing him to Caristi's theorem and for many fruitful discussions.

REFERENCES

- [1] J. Caristi, *Fixed point theorems for mappings satisfying inwardness conditions*, Transactions of the American Mathematical Society **215** (1976), 241–251.
- [2] D. Dzhafarov, S. Flood, R. Solomon, and L. Westrick, *Effectiveness for the dual Ramsey theorem*, Notre Dame J. Form. Log. **62** (2021), no. 3, 455–490.
- [3] I. Ekeland, *On the variational principle*, Journal of Mathematical Analysis and Applications **47** (1974), no. 2, 324–353.
- [4] D. Fernández-Duque, P. Shafer, and K. Yokoyama, *Ekeland's variational principle in weak and strong systems of arithmetic*, Selecta Mathematica **26** (2020), no. 68.
- [5] A. Freund, *What is effective transfinite recursion in reverse mathematics?*, MLQ Math. Log. Q. **66** (2020), no. 4, 479–483.
- [6] F. Khojasteh, E. Karapinar, and H. Khandani, *Some applications of Caristi's fixed point theorem in metric spaces*, Fixed Point Theory and Applications **16** (2016), no. 1.
- [7] W. A. Kirk, *Transfinite methods in metric fixed-point theory*, Abstract and Applied Analysis **2003** (2003), no. 5, 311–324.
- [8] A. Marcone, *On the logical strength of Nash-Williams' theorem on transfinite sequences*, Logic: from foundations to applications (Staffordshire, 1993), 1996, pp. 327–351.

- [9] W. Peng and T. Yamazaki, *Two kinds of fixed point theorems and reverse mathematics*, *Mathematical Logic Quarterly* **63** (2017), no. 5, 454–461.
- [10] S. Priess-Crampe and P. Ribenboim, *Ultrametric spaces and logic programming*, *The Journal of Logic Programming* **42** (2000), no. 2, 59–70.
- [11] ———, *Ultrametric dynamics*, *Illinois J. Math.* **55** (2011), no. 1, 287–303.
- [12] Y. Savateev and D. S. Shamkanov, *Non-well-founded proofs for the Grzegorzczuk modal logic*, *Rev. Symb. Log.* **14** (2021), no. 1, 22–50.
- [13] S. G. Simpson, *Subsystems of Second Order Arithmetic*, Second, *Perspectives in Logic*, Cambridge University Press, Cambridge; Association for Symbolic Logic, Poughkeepsie, NY, 2009.
- [14] H. Towsner, *Partial impredicativity in reverse mathematics*, *J. Symbolic Logic* **78** (2013), no. 2, 459–488.
- [15] X. Yu, *Riesz representation theorem, Borel measures and subsystems of second-order arithmetic*, *Ann. Pure Appl. Logic* **59** (1993), no. 1, 65–78.

DEPARTMENT OF MATHEMATICS WE16, GHENT UNIVERSITY, GHENT, BELGIUM AND INSTITUTE OF COMPUTER SCIENCE, CZECH ACADEMY OF SCIENCES, PRAGUE, CZECH REPUBLIC

Email address: fernandez@cs.cas.cz

URL: <https://users.ugent.be/~dfernnde/>

SCHOOL OF MATHEMATICS, UNIVERSITY OF LEEDS, LEEDS, UK

Email address: p.e.shafer@leeds.ac.uk

URL: <http://www1.maths.leeds.ac.uk/~matpsh/>

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF PENNSYLVANIA, PHILADELPHIA, USA

Email address: htowsner@math.upenn.edu

URL: <https://www.sas.upenn.edu/~htowsner/>

MATHEMATICAL INSTITUTE, TOHOKU UNIVERSITY, SENDAI, JAPAN

Email address: keita.yokoyama.c2@tohoku.ac.jp

URL: <http://www.math.tohoku.ac.jp/english/people/yokoyama-e.html>